

# Southern Hemisphere porbeagle shark (*Lamna nasus*) stock status assessment

## Covering Note on Stock Status and Conservation Advice (rev. 1)

Shelley Clarke  
Technical Coordinator-Sharks and Bycatch  
Common Oceans (ABNJ) Tuna Project  
WCPFC Secretariat, Pohnpei, Federated States of Micronesia

In recent years strong conservation and management actions have been taken for porbeagle sharks, mainly on the basis of evidence available for the Northern Hemisphere population in the Atlantic. The species as a whole was listed on the Convention on the Conservation of Migratory Species (CMS) in 2008 and on the Convention on International Trade in Endangered Species (CITES) Appendix II as of 2014. However, the status of the Southern Hemisphere stock has remained poorly understood in part because of the dispersed, circumpolar extent of its range and the absence of a mechanism to coordinate an assessment. This study attempts to assess the status of the Southern Hemisphere population of porbeagle shark (*Lamna nasus*) for the first time.

This study was conducted under the auspices of the Western and Central Pacific Fisheries Commission (WCPFC) and the Commission for the Conservation of Southern Bluefin Tuna (CCSBT) with funding from the Common Oceans (ABNJ) Tuna Project. The key data holders, which were identified as Argentina, Chile, Japan, New Zealand and Uruguay, had not previously collaborated on a stock status assessment and are not all members of WCPFC or CCSBT. As a result, rather than centrally compiling the data into a single dataset, this study was designed so that the consultant worked with each data holder to produce stock status indicators, and those indicators provided input to and context for a model-based assessment framework. Onsite analysis was undertaken in three of the five countries and webinars were used to share information amongst the study participants.

Initially it was proposed to use both a spatially-explicit risk assessment model, as well as a more traditional stock assessment model such as SS3, to assess the sustainability of fishery impacts on porbeagle sharks (see Section 2.7). Ultimately, however, the decision to split the study area into five sub-regions made it impractical to apply an age-structured model in the available time. One of the main benefits of an age-structured model would have been to address variation in selectivity, and subsequent effects on the population, as either the fishery, or the population distribution/size structure, changes. The likelihood of such change is considered to be minimal, however, it would be possible to investigate the selectivity issue through further development of the risk assessment model. This could shed more light on how much of the adult Southern Hemisphere porbeagle population inhabits what is effectively a refuge from fishing south of 45°S.

The risk assessment modelling results presented in this paper evaluate whether current impacts from fisheries exceed a maximum impact sustainable threshold (MIST, a kind of limit reference point) defined based on population productivity (see Figure 17). This approach differs from traditional stock assessments because it evaluates whether the population's ability to

withstand fishing pressure is exceeded, rather than evaluating biomass and whether the population is overfished. Therefore, this study examines the question of whether overfishing is occurring but does not answer questions regarding whether the stock is depleted. The age-structured model might have been able to address both overfishing and depletion but it is unlikely that the available data inputs would have allowed for robust estimation.

The risk assessment model estimated that fishing mortality on the Southern Hemisphere stock is very low, and that it decreases eastward from the waters off South Africa to the waters off New Zealand. For the assessment presented here, estimated values of fishing mortality were compared to a MIST ( $F_{\text{crash}}$ ) which indicates a level of fishing expected to lead to population extinction in the long-term. For all regions combined (Eastern Atlantic Ocean to Western Pacific Ocean) the fishing mortality was less than 10% of the MIST in all years assessed (1992-2014) and fell to half that level in more recent years, with at most a 3% probability of exceeding the MIST in 2010-2014 (see Section 2.6.4). This scenario is based on 100% capture mortality; assuming that some porbeagles survive their encounter with the fishery would reduce the estimated risk levels even further (see Section 3). For the alternative MISTs, also based on life history parameters, fishing mortality is less than 12% of the  $F_{\text{lim}}$  and less than 18% of the  $F_{\text{msm}}$  (see Addendum to this report). Further direction on which MIST/LRP should be applied will be incorporated into a final version of this assessment report.

The greatest contributions to fishing mortality were made by the pelagic longline fisheries, with the largest contribution (70-90%) from fleets targeting southern bluefin tuna (SBT) or a mixture of SBT and albacore tuna (ALB; see Figure 24). The contribution of the mixed ALB/SBT fishery increased considerably from about 2004 in the Western Indian/Eastern Atlantic Ocean region, and from 2005 in the Eastern Indian Ocean region, but has been consistently important in the Western Pacific Ocean region since 1996. In these regions, the more northerly longline fishery that takes albacore, bigeye and yellowfin tunas made a generally smaller but still substantial contribution to Southern Hemisphere porbeagle fishing mortality in recent years. In the Western Pacific region, the New Zealand midwater trawl fishery also contributed to this fishing mortality (see Section 2.6.5).

Although this study succeeded in obtaining a reasonable amount of data on the Southern Hemisphere porbeagle stock, there are many areas throughout the apparent range of this stock that remain distinctly data poor. In addition to improving data for the fisheries analysed here, it is also important to ensure that other fisheries catching Southern Hemisphere porbeagles are compiling catch records and collecting biological samples whenever possible (see Section 3). Publication of this study may present an opportunity to contact other regional fishery bodies, perhaps through the Common Oceans (ABNJ) Deep Seas Project, and advocate the importance of data improvement for the Southern Hemisphere porbeagle.

## Southern Hemisphere porbeagle shark stock status assessment

Prepared for Western and Central Pacific Fisheries Commission

S.D. Hoyle<sup>1</sup>, C.T.T. Edwards<sup>2</sup>, M.-J. Roux<sup>2</sup>, S.C. Clarke<sup>3</sup>, M.P. Francis<sup>2</sup>

<sup>1</sup> National Institute of Water and Atmospheric Research, P.O. Box 893, Nelson, New Zealand

<sup>2</sup> National Institute of Water and Atmospheric Research, Private Bag 14901, Wellington, New Zealand

<sup>3</sup> ABNJ (Common Oceans) Tuna Project, Western and Central Pacific Fisheries Commission, Pohnpei, Federated States of Micronesia, 96941

### Draft Final Report

This report will be revised based on comments received from reviewers and a final report prepared for WCPFC by the end of September 2017

July 2017

## Contents

|                                                                                |           |
|--------------------------------------------------------------------------------|-----------|
| <b>Executive summary .....</b>                                                 | <b>3</b>  |
| <b>1     Introduction .....</b>                                                | <b>6</b>  |
| 1.1   Historical background .....                                              | 6         |
| 1.2   Biology and distribution.....                                            | 7         |
| 1.3   Population trends .....                                                  | 9         |
| 1.4   Current conservation and management designations and measures.....       | 13        |
| 1.5   Status evaluation .....                                                  | 14        |
| <b>2     Methods and Results .....</b>                                         | <b>16</b> |
| 2.1   Assessment structure .....                                               | 16        |
| 2.2   Effort data .....                                                        | 17        |
| 2.3   Population distribution / density .....                                  | 18        |
| 2.4   Catch data and estimation .....                                          | 23        |
| 2.5   Indicator analyses .....                                                 | 27        |
| 2.6   Risk assessment .....                                                    | 32        |
| 2.7   Quantitative stock assessment .....                                      | 46        |
| <b>3     Discussion .....</b>                                                  | <b>46</b> |
| <b>4     Acknowledgements .....</b>                                            | <b>49</b> |
| <b>5     References.....</b>                                                   | <b>50</b> |
| <b>Appendix A             R code for calculating areas of grid cells .....</b> | <b>56</b> |
| <b>Appendix B             Fishing mortality estimates.....</b>                 | <b>57</b> |
| <b>Appendix C             F-ratios .....</b>                                   | <b>60</b> |

## Executive summary

This report presents the results of a Southern Hemisphere status assessment of porbeagle shark. The study was a collaborative one involving many countries with Southern Hemisphere fisheries that catch porbeagles. Participating scientists from Argentina, Chile, Japan, New Zealand and Uruguay contributed data analyses and abundance indices. Our approach combined indicator analyses and a spatially-explicit sustainability risk assessment. Indicator analyses were performed independently for different Southern Hemisphere fisheries and served to characterise local trends in relative abundance based on commercial catch per unit effort (CPUE) data, and trends in size and sex ratio based on biological data.

We limited our analyses to the region south of 30 °S which provided most of the available data, although the porbeagle shark's range extends slightly north of this latitude. Porbeagle sharks are taken in fisheries at least as far south as 56 °S. Southern Hemisphere population structure is not well understood, and we considered it unlikely that the population comprises a single well-mixed stock for management purposes. We subdivided the spatial domain of the assessment into five subpopulations or regions by longitude: 1) Western Atlantic Ocean; 2) Western Indian/Eastern Atlantic Ocean; 3) Eastern Indian Ocean; 4) Western Pacific Ocean; and 5) Eastern Pacific Ocean.

We applied different assessment methods by region, depending on data availability and quality. In the Western Indian/Eastern Atlantic Ocean, Eastern Indian Ocean, and Western Pacific regions, stock status assessment was performed using a spatially-explicit risk assessment. Indicator-based analyses were used to assess stock condition in the Eastern Pacific and the Western Atlantic, where there was limited information. We compared results from areas with varying levels of information, for greater insight into the status of the stock, levels of uncertainty, and data requirements for future studies.

Public domain surface longline data were obtained at a resolution of 5 x 5° grid by month by flag from regional fishery management organisations. Catch and effort data were also obtained from other trawl and longline fisheries known to take porbeagle sharks. Japanese observer data on catch and effort throughout the Southern Hemisphere were analysed to determine relationships between catch rates and the covariates year, quarter, latitude, hooks between floats, hooks, and sea surface temperature. These relationships were then used to predict relative abundance across the entire spatial domain, and combined with effort to predict surface longline catches. Catch estimates for other fisheries were obtained from the literature.

Most catch rate indicators were relatively short, variable, and uncertain, with the majority either stable or increasing. Length indicators were also variable. Only the Argentinian size and sex indicators showed temporal trends, with a small decline in sizes for both sexes, and a slight trend towards less female bias in the sex ratio index.

The indicator analyses, in addition to providing time series to monitor population change, revealed spatial patterns in size and sex distributions, and relationships with environmental variables. Such analyses are critical inputs to stock status assessments, because they help to determine model structure.

The risk assessment uses a quantitative framework to estimate spatially-explicit fishing mortality. It derives sustainability status as the ratio of total impact to a maximum impact sustainable threshold (MIST) reference point. The quantitative framework quantifies and propagates uncertainty throughout the assessment process. The risk assessment served to integrate selected CPUE

indicators in the evaluation of risk from commercial pelagic longline fisheries to porbeagle shark within an area subset of the Southern Hemisphere. The spatial domain of the risk assessment covered three regions: Western Indian/Eastern Atlantic Oceans, Eastern Indian Ocean, and Western Pacific Ocean, bounded at 30 °S and 60 °S. The Western Indian/Eastern Atlantic Ocean region was selected as the 'calibration region', being the most data-rich. A biomass dynamic model was fitted to the estimated catch and the abundance index for the calibration area. The model estimated a catchability parameter for the pelagic longline effort, which was used to estimate fishing mortality for the calibration area, and extended to other model areas.

Annual fishing mortalities were greatest in the Western Indian/Eastern Atlantic Ocean, slightly lower in the Eastern Indian Ocean, and lowest in the Western Pacific Ocean. Annual median  $F$  decreased from the mid-1980s to 2014 in both the Western Indian Ocean/Eastern Atlantic Ocean and Eastern Indian Ocean regions. In the assessment area (three regions combined) in the last decade (2005 to 2014), median  $F$  values ranged from 0.0008 to 0.0015 (mean 0.0010).

Risk was determined from the relationship between impact and the MIST limit reference point for the stock, which is the fishing mortality that will lead to a population crash. Risk values are calculated both as an  $F$ -ratio (Impact / MIST) and the probability that  $F$  exceeds the MIST, for the period from 1992 onwards (the first year of Japanese CPUE data).  $F$ -ratios for the assessment area declined by half from a mean of 0.068 (range 0.051—0.088) in 1992-2005, to a mean of 0.032 (range 0.023—0.042) in 2006-2014. The probability of  $F$  exceeding the MIST decreased by 95% from a mean of 0.0084 (range 0.0015—0.0205) in 1992-2005, to a mean of 0.0004 (range 0.0000—0.0013) in 2006-2014.

In the last 10 years, the SBT and ALB/SBT fisheries combined contributed about 75-80% of the fishing mortality in the Western Indian Ocean/Eastern Atlantic Ocean, 70-90% in the Eastern Indian Ocean, and 70-85% in the Western Pacific Ocean

Thus, results from the risk assessment indicate low fishing mortality rates in the three regions comprising the assessment area, and low risk from commercial pelagic longline fisheries to porbeagle shark over the spatial domain of the assessment. These results are consistent with the trends observed in catch rate indicators over the entire Southern Hemisphere range of the porbeagle shark population, which in most cases show stable or increasing catch rates. Concern has previously been expressed about reduced catch rates in the Uruguay longline fishery after 1993, but this concern is allayed by the re-analysis undertaken in collaboration with this project.

The population catchability was calibrated assuming that capture mortality was 100% (i.e., post-release survival is zero). Allowing for post-release survival would reduce these fishing mortality estimates, and reduce the estimated risk.

The catch rate indicators are the most important factors driving the results of the status assessment, and their reliability determines its reliability. The indicator trend in the calibration area is the most important factor determining the relatively low estimate of risk.

The risk assessment assumes that population density from 45 to 55 °S is the same as at 40 to 45 °S, since we have evidence from fisheries and surveys that they occur at these latitudes, but do not have Japanese longline observer data with which to estimate density. This is an important assumption, because it implies that the low fishing effort south of 45 °S provides a refuge from fishing mortality for the population. Biological data, and estimated relationships between size and SST, suggest that a high proportion of the adult population occurs at these latitudes.

Continued data collection by observers will improve the time series and provide better evidence about abundance trends. Maintaining collection and analysis of indicators from observer data is a key recommendation from this project. If further work is required, the following analyses could be carried out with currently available data:

- Explore assumptions about population density distribution and their effects on risk estimates, by rerunning the assessment with alternative density estimates.
- Explore selectivity at age in the Japanese pelagic longline data, which may permit estimation of the availability at age of the population to fishing. This analysis may permit two further developments: an age-structured analogue of the biomass dynamic risk assessment; and direct estimation of the proportion of the population south of 45 °S, removing the need to assume constant density from 45 to 55 °S.
- Further explore available biological data, to understand why patterns differ among areas. For example, it would be useful to model the effects of SST on size and sex patterns in the Chilean swordfish fishery.

The following recommendation would require further data collection:

- Compile biological and catch rate data from fisheries occurring south of 45 °S, such as the Chilean demersal longline fishery. Some data from this fishery are currently available, and data collection is ongoing.

The following recommendation would require additional, separate studies:

- Study porbeagle distribution using various types of markers (genetics, microchemistry, stable isotopes, parasites, conventional and electronic tags) to identify biologically-based boundaries.

The multiple indicators/risk assessment approach used in this study served to 1) source and synthesise available information on porbeagle shark at the scale of the Southern Hemisphere; 2) identify important data gaps (e.g., density distribution and life-stage specific vulnerability and overlap with fishing activities); 3) define a productivity-based reference point for the species; and 4) prioritise fishery areas for monitoring and management. This project has filled important information gaps by both directly analysing available life history information, and providing statistical support to the analyses by participating national fisheries scientists.

The project has provided the first assessment of the sustainability of the impact of fishing on the Southern Hemisphere porbeagle shark stock, and laid a foundation for future work. Results indicate that the impact of fishing is low across the entire Southern Hemisphere range of the porbeagle shark population.

# 1 Introduction

## 1.1 Historical background

The Western and Central Pacific Fisheries Commission (WCPFC) is one of five tuna Regional Fisheries Management Organisations (t-RFMOs) responsible for the sustainable use, conservation and management of highly migratory species taken by tuna fisheries. Unlike some of the other t-RFMOs, the WCPFC has explicit responsibility for assessing and managing not only tuna species, but also dependent and associated species under Articles 5(d) and 10.1(c) of its Convention. Recognition by the WCPFC of sharks as dependent and associated species in need of conservation and management has resulted in a list of fourteen shark species found in the Western and Central Pacific Ocean (WCPO) for which both data provision and assessment are required (Western and Central Pacific Fisheries Commission 2012). The WCPFC designated the porbeagle shark (*Lamna nasus*) as a key species at its seventh annual meeting in December 2010 but only in areas south of 20 °S due to concerns about species mis-identification in more northerly areas.

The designation of porbeagle as a key species by WCPFC may have been motivated by conservation and management proposals under other intergovernmental treaty organisations (see below) rather than by specific threats posed by WCPFC fisheries per se. This is because none of the WCPFC purse seine effort and only 7% of WCPFC longline fishing effort lies below 20 °S (based on SPC's Catch Effort Query system's raised aggregate data for 2013-2015 for longline effort, and Williams & Terawasi (2016)). As identified in a recent analysis of key shark species conducted by WCPFC's Scientific Services Provider, the Pacific Community (SPC), porbeagle sharks have been recorded in WCPFC observer datasets only in or immediately adjacent to the Australian and New Zealand Exclusive Economic Zones (EEZs; Rice et al. 2015), which represent only a small portion of the range of the Southern Hemisphere porbeagle stock. For this reason, while WCPFC committed to assessing the porbeagle shark's stock status by designating it as a key shark species, it was recognised that a broader regional approach would be necessary to undertake a comprehensive assessment (Rice et al. 2015).

At approximately the same time (March 2012), the Convention for the Conservation of Southern Bluefin Tuna (CCSBT) also identified porbeagle shark as a species of interest. After New Zealand compiled metadata on porbeagle biology, life history and catch and effort data to support an assessment in 2013 (Clarke et al. 2013) the CCSBT Ecologically-Related Species Working Group (ERSWG) in March 2015 agreed to request the Common Oceans (Areas Beyond National Jurisdiction (ABNJ)) Tuna Project, through its Technical Coordinator-Sharks and Bycatch position based at the WCPFC, to progress this work with the ERSWG and across the joint t-RFMOs. The ERSWG made this request on the basis that it would facilitate access to a broader range of data sets than would be available through the ERSWG members alone, and importantly cover the whole stock for assessment.

The Common Oceans (ABNJ) Tuna Project ([www.commonoceans.org](http://www.commonoceans.org)) is a partnership between the five t-RFMOs, as well as governments, inter- and non- governmental organisations, and the private sector, aimed at sustainable and efficient tuna fisheries production and biodiversity conservation. It focuses its efforts on marine resources that do not fall under the responsibility of any one country, thus working both in coastal and high seas areas. One set of activities of this GEF-funded project aims at reducing the impact of tuna fisheries on biodiversity by improving data and assessment methods for sharks, thereby promoting their sustainable management. Within this set of activities WCPFC is leading four stock status assessment studies for Pacific-wide shark stocks. The first study, a stock



status assessment of the bigeye thresher shark (*Alopias superciliosus*) was completed in September 2016. The Southern Hemisphere porbeagle shark was selected as another species of interest as its distribution is not only pan-Pacific but global, making a cooperative, inter-regional approach particularly important. The objectives of the Common Oceans (ABNJ) Tuna Project assessments include evaluating whether current t-RFMO management schemes are adequate, supporting national management actions such as Convention on International Trade in Endangered Species (CITES) Non-Detriment Findings, and demonstrating new modes of international cooperation for the assessment of highly migratory species.

## 1.2 Biology and distribution

Porbeagle sharks are cold-temperate, wide-ranging, coastal and oceanic sharks (Compagno 2001). Recent studies show they undergo both diel and reverse diel vertical movement patterns, and exhibit coastal site fidelity as well as large-scale open ocean movement (Pade et al. 2009, Campana et al. 2010b, Saunders et al. 2011, Francis et al. 2015). This species is distributed in the Northern Hemisphere from approximately 20°–75 °N but only in the Atlantic Ocean and Mediterranean Sea. It is absent from the North Pacific Ocean, where the closely related salmon shark, *Lamna ditropis*, fills its niche. In contrast, Southern Hemisphere porbeagle sharks have a circumpolar distribution (Last & Stevens 2009, Ebert et al. 2013). Although they are mainly caught between 30 °S and 50 °S, in the South Pacific porbeagles have sometimes been caught further north in the austral winter (June to August) and spring (September to November); in summer (December to February), they are not found north of about 35 °S. In summer and autumn, Southern Hemisphere porbeagle sharks appear to penetrate further south, and are found near many of the sub-Antarctic islands in the Indian and South-west Pacific Oceans (Francis & Stevens 2000).

The Northern and Southern hemisphere porbeagle shark populations are genetically and biologically distinct, and geographically disjunct (Figure 1; Testerman et al. 2007, Kitamura & Matsunaga 2010). As a result, the two populations have quite different life history characteristics: the Southern Hemisphere porbeagle is a smaller form that grows more slowly and lives twice as long as its northern conspecifics (Francis et al. 2007, Clarke et al. 2015). In the North Atlantic, porbeagles are often found close to shore but they also occur in the open ocean: mature females make long migrations into the subtropical waters of the central North Atlantic to give birth (Pade et al. 2009, Campana et al. 2010b, Saunders et al. 2011, Biais et al. 2017). In the Southern Ocean porbeagles are commonly caught in pelagic habitats far from shore, but also occur in coastal waters (Yatsu 1995, Francis & Stevens 2000, Semba et al. 2013, Francis et al. 2015). Limited tagging results from New Zealand confirm that Southern Hemisphere porbeagles undergo seasonal north-south movements and some make longitudinal movements of several thousand kilometres (Francis et al. 2015). It is not known whether there is a single circumpolar population in the Southern Hemisphere or whether there are multiple stocks or sub-stocks spread over this wide range.

Life history data for the Southern Hemisphere population derive primarily from studies in New Zealand and Australia; there is scant life history information from other Southern Hemisphere areas (International Commission for the Conservation of Atlantic Tunas 2010, Clarke et al. 2015). Length at birth is 58–67 cm fork length (FL; Francis & Stevens 2000); females mature at around 170–180 cm FL (age 13–16 years) and males at about 140–150 cm FL (age 6–8 years) (Francis & Duffy 2005, Francis 2015). Longevity is unknown but may be more than 65 years (Francis et al. 2007). Porbeagles are live-bearers (aplacental viviparous) and exhibit uterine oophagy with embryos feeding on other ova produced by the mother (Francis & Stevens 2000). The gestation period is about 8–9 months. Litter size is usually four embryos, with a mean litter size in the South-West Pacific of 3.75 (Francis &

Stevens 2000). If the reproductive cycle lasts one year, annual fecundity would be about 3.7 pups per female (Francis et al. 2008).




Figure 1. World distribution of porbeagle shark. NB: The northern and southern range limits of the Southern Hemisphere population are not well known and may be unreliable. Source: IUCN (<http://maps.iucnredlist.org/map.html?id=11200>).

Two studies of the age and growth of New Zealand porbeagles produced growth curves that show that males and females grow at similar rates up to about 10 years of age (about 150cm FL) and diverge thereafter with male growth approaching an asymptote while females continue to grow at a similar rate (Figure 2). The second study (Francis 2015) obtained slightly younger ages for a given length than the earlier study (Francis et al. 2007) because of a modified vertebral band pair counting protocol. Both studies cautioned that growth parameters are probably only accurate for ages up to about 20 years (because growth bands in older sharks become too narrow to be resolvable with a light microscope) and require further validation (Clarke et al. 2015).

In New Zealand, porbeagle sharks recruit to commercial fisheries during their first year at about 70 cm FL, and much of the commercial catch is immature (Francis 2015). Most sharks caught by tuna longliners are 70-170 cm FL. The size and sex distribution of both sexes is similar up to about 150 cm, but larger individuals caught by New Zealand fisheries are predominantly male; few mature females are caught. Regional differences in length composition suggest segregation by size (Francis 2013). The size and sex composition of sharks caught by trawlers are unknown. Porbeagles are active pelagic predators mainly of fish, but squid are also commonly eaten especially by the small sharks (Griggs et al. 2007, Horn et al. 2013).




Figure 2. Growth curves for porbeagle shark (reproduced from Clarke et al. 2015).

A study of the intrinsic rate of population increase of 38 species of sharks indicated that porbeagle shark has low productivity, similar in reproductive potential to some of the coastal carcharhinids (such as *Carcharhinus plumbeus* (sandbar shark) and *C. obscurus* (dusky shark)) and the pelagic thresher shark (*Alopias pelagicus*) (Cortés 2002). A comparison of productivities of twenty pelagic shark stocks in the Atlantic suggested that the porbeagle shark has relatively lower productivity than most species examined (13th of 20), although it had higher productivity than the bigeye thresher (*Alopias superciliosus*) and higher or similar to mako sharks (*Lamna* spp.) (Cortés et al. 2015). In terms of overall vulnerability (i.e. productivity and susceptibility) the porbeagle was the third most vulnerable shark of the 20. It should be noted, however that these studies were based on life history characteristics of the Northern Hemisphere porbeagle population. A similar ecological risk assessment of the Southern Hemisphere population in the Indian Ocean suggested the porbeagle was the seventh most vulnerable of the 17 species considered (Murua et al. 2012). Given that the Southern Hemisphere porbeagle has a longer generation time than its Northern Hemisphere conspecific, it may be more vulnerable to depletion.

### 1.3 Population trends

To date, most population-level studies of porbeagle sharks have been conducted for the North Atlantic, where this species has been highly valued for its meat and as a popular target for recreational fishing for several decades. As a result, porbeagle stocks in the North Atlantic currently show signs of serious overfishing in the form of greatly diminished catches compared to peak periods (Campana et al. 2008, Food and Agriculture Organization of the United Nations 2017). There has

been less attention to the status of Southern Hemisphere stocks, perhaps in part owing to catches from the southern stock being generally incomplete (Food and Agriculture Organization of the United Nations 2017). While there is no known, thriving market for its meat in the Southern Hemisphere, porbeagles have in the past been utilized for their fins (e.g. in the southern bluefin tuna fishery) as well as retained whole in countries such as New Zealand (Clarke et al. 2013, Food and Agriculture Organization of the United Nations 2017).

The most comprehensive attempt to assess porbeagle stocks was conducted by the International Commission for the Conservation of Atlantic Tunas (ICCAT) in 2009 (International Commission for the Conservation of Atlantic Tunas 2010). Separate analyses were conducted for the northeast, northwest and South Atlantic (Figure 3).

In the northeast Atlantic, there was considerable uncertainty in identifying the current stock status relative to virgin biomass because the peak of the fishery occurred well before the earliest points in the abundance indices. Nevertheless, the ICCAT assessment agreed with the view of the Northeast Atlantic Fisheries Commission (NEAFC, the Regional Fishery Body) that the stock was in a depleted state. It found that if catches were limited to zero, the stock would rebuild to its maximum sustainable yield level in 15-34 years, but if the current allowable catch was maintained rebuilding would require longer, possibly over 100 years (International Commission for the Conservation of Atlantic Tunas 2010).




Figure 3. Catch per unit effort series for the northwest Atlantic (upper figures), northeast Atlantic (lower left figures) and southwest Atlantic (lower right figures) stocks (ICCAT 2010).

In the northwest Atlantic, ICCAT's work was compared to, and found to agree with, an earlier Canadian stock assessment for coastal waters (Campana et al. 2010a). Both assessments concluded that the population is highly depleted but recovering under current management implemented by Canada and the United States. However, depending on the stock productivity and fishing mortality assumptions applied, recovery is projected to be achieved on the order of decades to over 100 years (International Commission for the Conservation of Atlantic Tunas 2010, Campana et al. 2012). There was no appreciable difference in the results when previously unaccounted for high seas catches were estimated and included in the model (International Commission for the Conservation of Atlantic Tunas 2010).

ICCAT's assessment for the South Atlantic was hampered by limited data for southwest and southeast stocks. In the southwest there was an apparent decline in catch rates in the Uruguayan fleet, with models suggesting that overfishing was occurring and that the stock was overfished. In the southeast, catch rates appeared stable since the early 1990s but biomass levels could not be estimated. The overall result for both stocks was that a robust conclusion on stock status could not be drawn (International Commission for the Conservation of Atlantic Tunas 2010).

A study based on a large Japanese observer and research survey dataset from 1982–2011 provided the most comprehensive view of the Southern Hemisphere stock (Figure 4) (Semba et al. 2013). One of the findings of that study was that large adult porbeagles penetrate into colder waters at higher latitudes, beyond the range of the southern bluefin tuna longline fishery, so they may not be subject to large-scale fishing pressure. In support of this theory the authors reported no declining trend in relative abundance in the Southern Ocean longline fishery from 1994–2011 (Semba et al. 2013).

In the Pacific, New Zealand has conducted indicators analyses for longline-caught porbeagle shark assessing trends in distribution, catch composition, abundance, size and sex ratios (Francis et al. 2014, Francis & Large 2017). There was some inconsistency among trends identified for porbeagle shark by the distribution and CPUE indicators, and by the standardised CPUE indices for the northern and southern New Zealand fisheries. Furthermore, some CPUE models fitted the data poorly and may be unreliable. Nevertheless, when taken as a group, the indicators suggested that the porbeagle population around New Zealand has been stable or increasing since 2005. Prior to that time observer data suggested a decline in abundance in the late 1990s and early 2000s followed by stability at a relatively low level.

SPC also conducted an indicators analysis which included porbeagle shark data for the wider South Pacific (Rice et al. 2015). That analysis benefitted from the inclusion of older records off Tasmania from the Australian observer programme in the 1990s (possibly identified as makos before then (Bruce 2014)), but mainly relied on the New Zealand observer records in more recent years (Figure 5). Not surprisingly, Rice et al. (2015) found the same pattern of high but variable porbeagle catch rates in the late 1990s followed by a low, fluctuating and slightly increasing catch rates thereafter. Rice et al. (2015) also concluded that most observed porbeagles were smaller than the size at maturity.




Figure 4. Catch per unit effort for porbeagle longline (top) and drift net (bottom) fishing gear in the Southern Ocean. In the top panel observer data are shown in red and survey data are shown in blue. Crosses denote no catch. (Semba et al. 2013).




Figure 5. Spatial distribution of the proportion of longline sets for which one or more porbeagle sharks were caught for each five-year period between 1995 and 2014 (Rice et al. 2015).

## 1.4 Current conservation and management designations and measures

The IUCN Red List classifies porbeagle sharks as “Vulnerable” based on population trends from the Northern Hemisphere and Uruguay as of 2006 (Stevens et al. 2006). Since that assessment several international organisations have adopted protections for the porbeagle as follows:

- 2008 Listed on Appendix II of the Convention on the Conservation of Migratory Species of Wild Animals (CMS), which encourages international cooperation toward conservation.
- 2010 Added to the CMS Memorandum of Understanding (MOU) for Sharks, which will develop a Conservation Plan to guide cooperation between the signatories to CMS Convention as well as other interested stakeholders (Convention on Migratory Species 2016)
- 2011 NEAFC, citing the porbeagle shark’s low productivity and high vulnerability to overfishing, prohibited directed fishing and mandated prompt release. This measure will remain in effect until the end of 2019 (Northeast Atlantic Fisheries Commission 2016).
- 2012 Added to Annex II of the Barcelona Convention. In response, the General Fisheries Commission for the Mediterranean (GFCM) agreed under GFCM/36/2012/3 to prohibit retention on board, trans-shipping, landing, transferring, storing, selling or displaying or offering for sale porbeagle specimens caught in the Mediterranean (Food and Agriculture Organization of the United Nations 2012).
- 2014 Listed on Appendix II of CITES, requiring that all exports of porbeagle sharks, including landings in non-flag State ports, be accompanied by permits issued by the flag state CITES Management Authority. Export permits are contingent upon legal acquisition and non-detriment findings (NDFs), the latter of which represents a certification by an authorized CITES Scientific Authority that the proposed export is not detrimental to the survival of the species (Clarke et al. 2014).
- 2015 ICCAT adopted a recommendation requiring that porbeagles be released promptly and unharmed, to the extent practicable (International Commission for the Conservation of Atlantic Tunas 2015).



Various countries also have adopted management measures for porbeagle specifically including:

|                              |                                                                                                                                                                                                       |
|------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| European Union               | Zero total allowable catch by European Union vessels (European Union 2015).                                                                                                                           |
| Canada and the United States | Catch limits for the northwest Atlantic stock (International Commission for the Conservation of Atlantic Tunas 2010, National Oceanic and Atmospheric Administration 2016).                           |
| New Zealand                  | Catch limits under a quota management system (Ministry for Primary Industries 2016).                                                                                                                  |
| Australia                    | A requirement to release porbeagles brought up alive in Australia (Australian Fisheries Management Authority 2017).                                                                                   |
| Uruguay                      | Since January 2013, a prohibition on retaining porbeagles by Uruguayan-flagged vessels and foreign vessels fishing in the Uruguayan EEZ fisheries (R. Forselledo, personal communication, July 2017). |
| Argentina                    | A requirement to release live porbeagles (and other shark species) longer than 1.6 m (F. Cortés, personal communication, July 2017).                                                                  |

Other organisations with fishing grounds lying within the range of the Southern Hemisphere porbeagle population (assumed to be south of 20 °S) have enacted measures applicable to sharks in general. No-retention measures for all commercial take of sharks have been adopted by the French Overseas Territories of New Caledonia and French Polynesia, the British Overseas Territory of Pitcairn, and the Cook Islands. The Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) implemented a moratorium on all directed shark fishing in the Antarctic region in 2006 and encourages the live release of incidentally caught sharks (Commission for the Conservation of Antarctic and Marine Living Resources 2006).

In response to a petition to list the porbeagle under the United States' Endangered Species Act, the National Oceanic and Atmospheric Administration (NOAA) undertook a comprehensive status review of both the Northern and Southern hemisphere populations in 2016. This resulted in a finding that neither population is currently in danger of extinction throughout all or a significant portion of its range or likely to become so in the foreseeable future and thus listing was not warranted. With regard to the Southern Hemisphere population in particular, NOAA's review found that abundance is stable or increasing, but that there is some uncertainty about current stock status (Curtis et al. 2016, National Oceanic and Atmospheric Administration 2016).

## 1.5 Status evaluation

This report presents the results of a Southern Hemisphere status assessment of porbeagle shark. The study was a collaborative one involving many countries with Southern Hemisphere fisheries that catch porbeagles. Participating scientists from Argentina, Chile, Japan, New Zealand and Uruguay contributed data analyses and abundance indices used and considered during the risk assessment. The study team supported this work by providing analytical advice to participating scientists during the development of the indicators.



The indicators are described in separate papers, which have been submitted to WCPFC-SC13 as Information Papers. Analyses of Japanese longline data (Hoyle et al. 2017b) provided catch rate, size and sex indicators for the Western Indian/Eastern Atlantic Ocean, Eastern Indian Ocean, and Western Pacific regions; a basis for estimating the spatial distribution of porbeagle sharks worldwide; and a basis for imputing catches in pelagic longline fisheries. Analyses of New Zealand fisheries data provided catch rate, size, and sex indicators (Francis & Large 2017) and catch estimates (Francis 2017) for the Western Pacific region. Analyses of Chilean swordfish fishery data provided catch rate, size, and sex indicators and catch estimates for the Eastern Pacific region (Hoyle et al. 2017a). Analyses of Uruguayan longline data provided catch rate indicators (Forselledo et al. 2017) for the Western Atlantic region. Analyses of Argentinean surimi trawl fishery data (Cortés et al. 2017) provided catch rate, size, and sex indicators for the Western Atlantic region.

Our approach combined indicator analyses and a spatially-explicit sustainability risk assessment. Indicator analyses were performed independently for different Southern Hemisphere fisheries/study partners and served to characterise local trends in relative abundance based on commercial catch per unit effort (CPUE) data, and trends in size and sex ratio based on biological data. We also considered a more complex age and length-structured assessment, using Stock Synthesis software (Methot & Wetzel 2013).

Risk assessment tools have been developed in response to data limitation problems in the evaluation of fishing effects on non-target species, including sharks and other elasmobranch species (Stobutzki et al. 2002, Braccini et al. 2006, Griffiths et al. 2006, Cortés 2008, Zhou & Griffiths 2008, Cortés et al. 2010, Gallagher et al. 2012, Cortés et al. 2015). We adapted and modified the risk assessment methods developed by Fu et al. (2016) in a stock status assessment of bigeye thresher shark, *Alopias superciliosus*. The method uses a quantitative framework for estimating spatially-explicit fishing mortality and deriving a sustainability status for the species as the ratio of total impact to a maximum impact sustainable threshold (MIST) reference point. Rather than following a traditional stock assessment approach, which relies heavily on population processes that for sharks are often poorly understood, this spatially-explicit approach is based on species productivity, inferred distribution and data on the occurrence, characteristics and intensity of fishing. The quantitative framework allows uncertainty to be quantified and propagated throughout the assessment process. An important outcome is that impact, sustainability risk and uncertainty can be partitioned spatially and among fishery sectors, allowing more focused management. The risk assessment served to integrate selected CPUE indicators in the evaluation of risk from commercial pelagic longline fisheries to porbeagle shark within an area subset of the Southern Hemisphere having the greatest amount of data. Indicator-based analyses were then used to assess condition in the remainder of the Southern Hemisphere (see Section 2.1). This combined approach allowed us to integrate results from areas with varying levels of information, and to gain greater insight into the status of the stock, levels of uncertainty, and the data requirements for future studies.

## 2 Methods and Results

### 2.1 Assessment structure

This study covers the entire Southern Hemisphere porbeagle shark population. Porbeagles have been reported in fisheries or surveys circum-globally in the Southern Hemisphere, so all longitudes are included. Porbeagles are found as far north as 20 °S, though catch rates are very low north of 30 °S. Porbeagle catches have been reported from further north, including at the equator, but logbook data include reporting errors, and we considered these northernmost catches to be errors. We limited analyses, and therefore the assessment domain, to the area south of 30 °S to avoid problems fitting analyses to strata without any catch. In the Southern Hemisphere, most longline porbeagle catch is taken north of 45 °S, but this is probably due to the distribution of the southern bluefin tuna longline fishery rather than the distribution of porbeagle sharks.

Porbeagle sharks are also taken in fisheries at least as far south as 56 °S, such as the mackerel icefish and Patagonian toothfish trawl and longline fisheries in the Heard and McDonald Island EEZ, in New Zealand midwater trawl fisheries, in the Argentinian surimi trawl fishery, and in the Chilean demersal longline fishery. Porbeagle sharks were observed in the eastern Pacific to the southern limits of the JAMARC longline survey (60 °S) and the JAMARC gillnet survey (52.5 °S) (Yatsu 1995, Semba et al. 2013, Hoyle et al. 2017b). Porbeagle sharks may be found elsewhere, but few data are available.

We considered that Southern Hemisphere porbeagles are unlikely to comprise a single well-mixed stock for management purposes. Initial observations of trends in population indices from Japanese longline data (the most comprehensive dataset available) suggested that they may vary spatially (Hoyle et al. 2017b), although reanalysis of the Japanese observer data shows reasonably stable catch rates across three regions (see Section 2.5). Nevertheless, the spatial scale of the Southern Hemisphere is very large relative to observed longitudinal movement rates of porbeagles (Francis et al. 2015). Depletion of one longitudinal band may take considerable time to affect the population outside that area. Fisheries interactions suggest a higher incidence of juveniles in northern areas (Semba et al. 2013) and the majority of shark movements appear to be in the north–south direction (Francis et al. 2015), suggesting that mixing occurs across latitudes.

The potential for subdivision is apparent, but the population structure is not well understood. It is possible that there are subgroups within the population that have not been identified. There is some evidence for population structuring by age class and/or reproductive class longitude as well as by latitude, based on analyses of size and sex patterns in Japanese and Chilean observer data (Hoyle et al. 2017b, Hoyle et al. 2017a). Genetic analyses have even suggested the possibility of independent populations within the South Atlantic (Kitamura & Matsunaga 2010).

Based on this understanding of stock structure, we subdivided the Southern Hemisphere porbeagle stock into five subpopulations or regions by longitude, with the divisions based on variation in fishing effort and on geographical features (Figure 6). For comparison, Northern Hemisphere porbeagle sharks in the North Atlantic are managed as two separate stocks, one on each side of 42 °W, given low levels of population interchange, and evidence for site fidelity and homing behaviour (Biais et al. 2017). The five regions defined as notional boundaries of Southern Hemisphere porbeagle subpopulations in this study were:

1. Western Atlantic Ocean (70° to 10° W);
2. Western Indian/Eastern Atlantic Ocean (10° W to 70° E);

3. Eastern Indian Ocean (70° to 140° E);
4. Western Pacific Ocean (140° to 180° E); and
5. Eastern Pacific Ocean (180° E to 70° W).

We applied different assessment methods by region, depending on data availability and quality. In the Western Indian/Eastern Atlantic Ocean, Eastern Indian Ocean, and Western Pacific regions, stock status assessment was performed using a spatially-explicit risk assessment. Indicator-based analyses were used to assess condition in the Eastern Pacific and the Western Atlantic, where there was limited information.




Figure 6: Spatial subdivision of the Southern Hemisphere porbeagle population into five regions.

## 2.2 Effort data

Public domain surface longline data were obtained at a resolution of 5 x 5° grid by month from the following regional fishery management organisations: the Western and Central Pacific Fisheries Commission (WCPFC), the Inter-American Tropical Tuna Commission (IATTC), the International Commission for the Conservation of Atlantic Tunas (ICCAT), the Indian Ocean Tuna Commission (IOTC), and the Commission for the Conservation of Southern Bluefin Tuna (CCSBT).

Each dataset was adjusted to the same reference frame, with location marking the centre of the 5 x 5° grid, and longitudes 0–360°. The WCPFC, CCSBT and ICCAT datasets were affected by the three-vessel confidentiality rule, according to which data are only reported for time-region strata that include data from at least three fishing vessels.

To address data loss due to the three-vessel rule, public domain catch and effort data for the Western and Central Pacific Ocean were requested from the WCPFC for the period 2004–2014, in two formats: a) stratified by year, month, 5 x 5° grid (latitude and longitude), and flag (WCP\_FLAG), and b) stratified by year, quarter, and 5° latitude (WCP\_LAT). Both public domain datasets omit strata that include fewer than three vessels, to avoid potential identification (Western and Central Pacific Fisheries Commission 2007), which meant that more data were omitted from the less aggregated dataset WCP\_FLAG. However, WCP\_FLAG has higher spatial resolution. We therefore used the WCP\_LAT dataset to calculate a multiplier with which to scale up the effort in WCP\_FLAG for each year, quarter and latitude band to match the total effort in WCP\_LAT, while retaining the distribution

by 5° grid and month. Public domain catch and effort data were also obtained from the WPCFC website for the period 1950–2014, stratified by year, month, and 5 x 5° grid.

Atlantic Ocean effort data were obtained from the Task II catch and effort database (<https://www.iccat.int/en/accesingdb.htm>) for the period 1961–2014. For longline, these data are aggregated by flag, year, month and 5 x 5° grids. ICCAT avoids identifying individual vessels by omitting strata with observations from fewer than three vessels. For this dataset no information on total effort was available for scaling, so total effort was underestimated.

Effort data for parties reporting to the CCSBT were obtained from the public domain file [https://www.ccsbt.org/userfiles/file/data/CEData\\_Longline.xlsx](https://www.ccsbt.org/userfiles/file/data/CEData_Longline.xlsx) for the period 1965–2015. These data are aggregated by flag, year, month and 5 x 5° grids. The CCSBT data are known to be affected by the three-vessel rule but could not be adjusted, so total effort was underestimated.

Indian Ocean effort data were obtained from the IOTC website (<http://www.iotc.org/documents/ce-longline>) for the period 1960–2015. These data are aggregated by flag, year, month and 5 x 5° grids. The IOTC further aggregates data prior to release at a coarser resolution wherever there would otherwise be potential to identify individual vessels. Thus, all catch and effort data were included in the IOTC dataset. A small amount of IOTC effort was reported in days rather than hooks, and this was omitted.

Effort data from the Eastern Pacific were obtained from the IATTC website for the period 1963–2014, aggregated by flag, year, month and 5 x 5° grids. This dataset includes all the longline data exactly as provided by the countries.

A check of CCAMLR data holdings in November 2016 revealed a total of three reported captures (and subsequent release) under the generic code ‘sharks, skates and rays’ in bottom longline fisheries (S. Mormede, NIWA, pers. comm.). Based on this, no further data were requested from CCAMLR.

Trawl data were provided at a resolution of 5 x 5° grid and month by the South Pacific Regional Fisheries Management Organisation (SPRFMO). No porbeagle catch was reported in this fishery.

## 2.3 Population distribution / density

The spatially-explicit risk assessment methodology uses the spatial overlap of fishing effort and population density to derive a risk metric. This requires estimation of relative population density over the spatial domain of the assessment. Spatially-explicit porbeagle density was estimated at the same spatial resolution as the available effort data (i.e., in 5 x 5 degree grids). Population distribution was inferred from the spatial component of catch rates in the Japanese tuna longline fishery. When standardising catch and effort data to produce indices of abundance, the spatial representation of the models included both latitude and longitude. However, this approach only allows relative distribution to be estimated for longitudes and latitudes for which we have Japanese longline effort data. By removing longitude from the models and including sea surface temperature, we could use known values of sea surface temperature to predict relative abundance for all locations between 30 °S and 45 °S, circum-globally.

Analyses used a delta lognormal approach, first modelling the probability of nonzero catch, and then modelling the distribution of catch rates in the nonzero catches.

$$por \neq 0 \sim yr + qtr + s(lat, k = 10) + s(hbf, k = 5) + s(hooks, k = 10) + s(SST, k = 10)$$

$$\log\left(\frac{por}{hooks}\right) \sim yr + qtr + s(lat, k = 10) + s(hbf, k = 5) + s(SST, k = 10)$$

Latitude (*lat*), hooks between floats (*hbf*) and sea surface temperature (*SST*) were modelled as continuous variables. Year (*yr*), and quarter (*qtr*) were modelled as categorical variables. The term '*s*' refers to a one-dimensional thin-plate regression spline smooth.

Residuals for the positive component of the model indicated that all variables were statistically significant (Table 1), with reasonable fit to the data but some skewness (Figure 7). Results indicated strong relationships between *SST* and catch rates, particularly for the probability of nonzero catch, and relatively stable catch rates through time (Figures 8 and 9).




Figure 7: Residual distribution plots for the lognormal positive observer data analysis for catch prediction.

Table 1: ANOVA table for variables in the binomial and positive components of the delta lognormal standardisation model. For smooth variables degrees of freedom (DF) are the effective degrees of freedom calculated by the *mgcv* package.

| Model type | Data type   | Parameter | DF  | F     | p-value  |
|------------|-------------|-----------|-----|-------|----------|
| Binomial   | categorical | op_yr     | 22  | 292.0 | < 2e-16  |
| Binomial   | categorical | qtr       | 3   | 52.4  | 2.52E-11 |
| Binomial   | smooth      | s(lat)    | 5.1 | 52.5  | 2.32E-09 |
| Binomial   | smooth      | s(hbf)    | 2.6 | 54.8  | 1.27E-11 |
| Binomial   | smooth      | s(hooks)  | 7.9 | 68.2  | 2.95E-11 |
| Binomial   | smooth      | s(SST)    | 7.8 | 844.0 | < 2e-16  |
| Positive   | categorical | op_yr     | 22  | 14.0  | < 2e-16  |
| Positive   | categorical | qtr       | 3   | 12.4  | 4.22E-08 |
| Positive   | smooth      | s(lat)    | 8.6 | 16.0  | < 2e-16  |
| Positive   | smooth      | s(hbf)    | 3.2 | 6.3   | 9.29E-05 |
| Positive   | smooth      | s(SST)    | 8.5 | 19.1  | < 2e-16  |




Figure 8: Probability of nonzero catch predicted for pelagic longline effort, for values of each covariate with other values held fixed. Fixed values were latitude 40 °S, HBF 11, SST 12 °C, Year 1992, and Quarter 0.125 (first quarter).




Figure 9: Predicted catch rate for nonzero pelagic longline effort, for values of each covariate with other values held fixed. Fixed values were latitude 40 °S, HBF 11, SST 12 °C, Year 1992, and Quarter 0.125 (first quarter).



Catch rates were predicted for the available pelagic longline effort at stratification of 5 x 5 grid by month, for a standard year. The original analysis used set level data, and predictions were made for a standard set. Each set was assumed to use 3000 hooks, with 11 hooks between floats. SST was predicted for the month and 5 x 5 grid based on the CSIRO CARS 2009 Atlas of Regional Seas (Ridgway et al. 2002), which provides monthly predictions averaged across years. Patterns of sea surface temperature varied strongly by latitude, longitude, and month (Figure 10). Please note that month is used to derive SST, but is not itself a covariate. Each month was allocated to a quarter, which was a covariate. Separate predictions were made for the probabilities of nonzero catch, and the catch rates in positive sets. For each stratum, the two predictions were multiplied to give expected catch rate. Each monthly distribution was normalized to have a maximum of 1.

We assumed that variation in catch rate with temperature and latitude was associated with relative abundance rather than catchability, and predicted relative density in space by month (Figure 11).

By using the same predictive model to estimate the population distribution for the entire spatial domain, this approach assumes that population densities, depletion levels, and trends are similar in all regions. The only differences are caused by SST and latitude. It also assumes that the relationships of latitude and sea surface temperature with catch rate estimated from the Japanese longline data are applicable to all other pelagic longline effort. The resulting predictions suggest relatively consistent longitudinal gradients in porbeagle density within latitudinal bands over the Southern Hemisphere. This may or may not represent true population density, as porbeagle distribution may be patchy, with aggregation in some areas.

## 2.4 Catch data and estimation

For most of the pelagic longline effort, direct information on catches was not available. Reporting in Japanese logbooks of porbeagle catches was generally poor before 2008. We had little information about porbeagle catch rates for most other fleets. The best available information was derived from the Japanese observer data. Catches were therefore estimated from the expected catch rates estimated above (Section 2.3) and the observed effort (see Section 2.2). Catch rates were predicted as for the relative density prediction, but for all years rather than a standard year. Predictions could only be generated for years with catch rate estimates. For years without such estimates, the catch rates for 1992 were applied as a conservative assumption. Expected catches per stratum were estimated by multiplying the observed effort by the expected catch rate.

Porbeagle catch estimates for all fisheries in the New Zealand EEZ, including midwater trawl and longline fisheries, were provided by New Zealand (Francis 2017). Trawl effort and estimated porbeagle catches were provided for the Argentinian surimi fleet (Cortés et al. 2017).

Observer data for the Chilean swordfish fishery were provided for three sectors: industrial longline, artisanal longline, and artisanal gillnet (Hoyle et al. 2017a). Coverage for the artisanal sectors was reported to be 3% and for industrial longline 87%. Observed catches were scaled up to annual catch estimates by dividing the catch records by year and sector by the appropriate coverage rate.

Longline catches in the region of the Kerguelen and Crozet islands in the Southern Ocean were provided by the French Muséum National d'Histoire Naturelle, Paris (Guy Duhamel pers. comm.), at a resolution of year and FAO area. Catch data were reported by the Australian Bureau of Agricultural and Resource Economics and Sciences (Heather Chapman, pers. comm.) for the Heard and McDonald Island trawl fishery for 1996–97 to 2014–15, and for the Macquarie Island Patagonian toothfish longline fisheries for 2008–09 to 2014–15.

Combined catches per year by region are summarised in Figure 12.




Figure 10: Sea surface temperature patterns by month from CARS data. Yellow represents higher SST, and red lower SST.




Figure 11: Predicted population relative distribution by month from the abundance prediction model. For each month, the area with highest density is assigned relative density of 1. Yellow indicates higher density, and red lower density. White indicates no information.




Figure 12: Total catch in number of porbeagle sharks per year by region.

## 2.5 Indicator analyses

Abundance indices through time were required as inputs into the risk assessment, and to serve as indicators of population trend and condition. The abundance indicators reported here are based on fisheries that operated within each of the five areas, and were taken to be representative of temporal trends in abundance. This contrasts with the estimates of relative abundance in space, estimated in a broad-based analysis across the 190 degrees of longitude for which we had access to Japanese observer data, presented above in Section 2.3.

For some regions, where available, we also present indicators of trends in size and sex ratio.

### 2.5.1 Western Indian/Eastern Atlantic Ocean

A reanalysis of the Japanese data is presented here, using a modification of the method used by Hoyle et al. (2017b) in order to address problems observed with non-normal distributions of the residuals. Most of the methods used remain the same, and apart from some necessary background information, only the approaches that were changed are described here.

Abundance indices for the Western Indian/Eastern Atlantic Ocean, Eastern Indian Ocean, and Western Pacific Ocean were estimated from catch rates in Japanese longline fisheries (Hoyle et al. 2017b). Data were grouped into fishing strategies using cluster analysis of species composition in the logbook data. Observer data and logbook data were linked based on the vessel callsign and set date.

We fitted generalized additive models in R (R Core Team 2017) using the package *mgcv* (Wood 2011). The previous approach used categorical variables for 5 x 5° spatial grids with generalized linear models, and failed to provide estimates for grids in which all observations had zero catch. Using the spatial smoothing available in *mgcv* avoided this problem. The previous approach also fitted indices for separate northern and southern areas in each region, but we combined these to give a single set of indices for each region, with better statistical precision and fewer missing values. Analyses used a delta lognormal approach, first modelling the probability of nonzero catch, and then modelling the distribution of catch rates in the nonzero catches.

$$por \neq 0 \sim yr + qtr + te(lon, lat, k = c(7, 7)) + s(hbf, k = 5) + s(hooks, k = 10) + cl + s(SST, k = 5)$$

$$\log\left(\frac{por}{hooks}\right) \sim yr + qtr + te(lon, lat, k = c(7, 7)) + s(hbf, k = 5) + cl + s(SST, k = 5)$$

Latitude (*lat*), longitude (*lon*), hooks between floats (*hbf*) and sea surface temperature (*SST*) were modelled as continuous variables. Year (*yr*), quarter (*qtr*), and cluster (*cl*) were modelled as categorical variables. The term 's' refers to a one-dimensional thin-plate regression spline smooth, and 'te' to a two-dimensional tensor product smooth.

Residuals were more normally distributed (Figure 13) than those estimated in previous analyses (Hoyle et al. 2017b), which used categorical variables for spatial effects. Indices are variable and do not provide strong evidence of long-term trends (Figure 14).

Size and sex indicators are also available from this fishery (Figure 15).




Figure 13: Residual distribution plots for lognormal positive observer data analyses for the Western and Eastern Indian Ocean, and the Western Pacific Ocean.




Figure 14: CPUE indices for each region and contributing country. Sources: (Cortés et al. 2017, Forselledo et al. 2017, Francis & Large 2017, Hoyle et al. 2017b, Hoyle et al. 2017a).

### 2.5.2 Eastern Indian Ocean

Indices for the Eastern Indian Ocean region were estimated in the same way as those for the Western Indian/Eastern Atlantic Ocean (Figure 13).

Size and sex indicators are also available from this fishery (Figures 15 and 16).

### 2.5.3 Western Pacific Ocean

Two sets of indices were estimated for the Western Pacific. The first set was based on catch rates in the Japanese longline fishery, and was estimated in the same way as those for the Western Indian/Eastern Atlantic Ocean.

The second set was based on analyses of catch rates in New Zealand longline fisheries. They were reported by Francis & Large (2017) in an update of the results of a previous analysis (Francis et al. 2014). CPUE indices were provided for catches in the Japanese charter tuna longline fishery in southern New Zealand (the Japan South fishery), separately for both logbook data and observer data. There were also indices from observer data in northern New Zealand for both domestic and Japanese charter vessels combined. Indices are shown in Figure 13.

Size and sex indicators are also available from this fishery (Figures 15 and 16).

### 2.5.4 Eastern Pacific Ocean

Indices for the Eastern Pacific Ocean are based on catch rates in the Chilean swordfish fishery, which takes porbeagle sharks as bycatch, particularly in the southern portion of the fishery. The fishery comprises three components, the industrial longline, artisanal longline, and artisanal gillnet. The longline fishery data were combined and analysed to produce indices of abundance (Hoyle et al. 2017a) (Figure 13).

The Chilean index is relatively short and variable, reflecting the fact that data are sparse because porbeagles are only taken at the southern extreme of the swordfish fishery. There is no indication of a temporal trend in these indices.

Size and sex indicators are also available from this fishery (Figures 15 and 16).

### 2.5.5 Western Atlantic Ocean

Two sets of indices were estimated for the Western Atlantic. The first set was based on catch rates in the Uruguayan longline fishery (Forselledo et al. 2017). This was an update of a previous analysis (Pons & Domingo 2010). There were a number of changes from the 2010 analysis, the most important being breaking the index into two parts (1981–1991 and 1992–2012). Prior to 1992 all participants were large-scale freezer vessels using Japanese-style multifilament longlines (about 2000 hooks per set). From 1992 these vessels were replaced, mostly by small-scale fresh-fishing vessels using American-style monofilament longlines (about 900 hooks per set), and two vessels using Spanish-style multifilament longline. Porbeagle shark catch rates were much lower for vessels using the American style longlines.

Neither the early nor the late Uruguayan indices shows a clear trend through time. The later index has a very low catch rate and is also very variable.

The second set of indices was based on catch rates in the Argentinean surimi fishery (Figure 14). This trawl fishery has comparatively low catch rates, but provides a useful dataset. The fishery is further

south than others reported here, and takes relatively large porbeagles, with catch rates increasing further south. Size and sex indicators are also available from this fishery (Figures 15 and 16).

The Argentinian index is short, but appears to show an increasing trend through time. Indices of mean size appear to decline slightly for both males and females. There is also a slight trend toward a lower proportion of females in the sex ratio index.




Figure 15: Standardized predictions (Japan (JP) and Chile) and annual measurements (New Zealand (NZ) and Argentina) of lengths in the catch, by region and contributing country. Sources: (Cortés et al. 2017, Francis & Large 2017, Hoyle et al. 2017b, Hoyle et al. 2017a).






Figure 16: Standardized predictions (Japan (JP) and Chile) and annual measurements (New Zealand (NZ) and Argentina) of proportion female in the catch, by region and contributing country. Sources: (Cortés et al. 2017, Francis & Large 2017, Hoyle et al. 2017b, Hoyle et al. 2017a).

## 2.6 Risk assessment

The risk assessment methodology uses the spatial overlap of fishing effort and population density to derive a risk metric. This requires estimation of a catchability coefficient, which is achieved by fitting a logistic production model to available data in the most data-rich of the assessment regions. The catchability scalar is then applied to effort overlap in the other regions to estimate a fishing mortality. The sum of spatially-explicit, annual fishing mortality (annual impact) is compared to a maximum impact sustainable threshold (MIST), which is a limit reference point derived from the intrinsic rate of population growth. Risk is estimated from the ratio of annual impact to the MIST, and expresses the probability, given the uncertainty, that total impacts exceed the MIST.

A summary of data inputs, analytical methods and key parameters is presented in Figure 17.




Figure 17. Conceptual representation of data inputs, analytical methods and key parameters used in spatially-explicit risk assessment of the Southern Hemisphere porbeagle shark. BDM = Bayesian state-space biomass dynamics model. The dashed outline box represents analytical methods applied to a region subset of the available data.

We used the risk assessment model to evaluate the risk from commercial pelagic longline fisheries to porbeagle shark within a subarea of the Southern Hemisphere (hereinafter referred to as the spatial domain of the risk assessment). This area ranges from 10 °W to 180 °E longitude and from 30 to 60 °S latitude (see Figure 6) and corresponds to the region covered by the Japanese tuna longline fishery (Semba et al. 2013). The risk assessment was restricted to this area as it contained sufficient information to estimate key components of the risk assessment, namely the species distribution (population density) and population catchability. The quantitative risk approach assumes that true population abundance is unknown, but that catch and effort information from an area of highest abundance (and comparatively high data availability) can be used in conjunction with relative density estimates to calibrate a catchability parameter for the assessed population and estimate spatially-explicit relative fishing mortality and total impact from fisheries over the spatial domain of the assessment. Our approach assumes that porbeagle sharks are distributed over the spatial domain of

the assessment, comprised of three regions/subpopulations (potentially distinct biological stocks) that undergo limited or no mixing: 1) Western Indian/Eastern Atlantic Ocean; 2) Eastern Indian Ocean; and 3) Western Pacific Ocean (see Section 2.1). We estimated fishing mortality and calculated risk separately for each of the assessment regions, and across the whole spatial domain.

The risk assessment is spatially-explicit and quantitative and distinguishes impact and risk among fishery sectors (i.e., fleet components characterised by differences in operational practice and therefore, different catchability). Annual impacts were estimated over a spatial grid of 5° latitude x 5° longitude, corresponding to the spatial resolution of the catch and effort data available for assessment. The timeframe of the bdms assessment was the period of commercial effort (logsheet) data from 1960 to 2014, though for the risk assessment metrics we give greater weight to the period with better data, starting in 1992. The fishing effort data were modelled as a single fleet (with hooks per set standardised to that of a standard vessel from the Japanese fleet). Only seasonal (year-quarter) variability in species distribution and operational practice (e.g., gear type and targeting strategies) were considered in the assessment. Uncertainties in species distribution within season were not considered. Uncertainty in the catchability parameter and population productivity parameter were estimated and propagated into the evaluation of risk.

### 2.6.1 Preparation of spatial data

The spatial domain of the assessment covered three regions: Western Indian/Eastern Atlantic Oceans, Eastern Indian Ocean, and Western Pacific Ocean, bounded at 30 °S and 60 °S (Figure 18). The Western Indian/Eastern Atlantic Ocean region is also referred to as the ‘calibration region’, being the most data-rich. Each region was divided into 5 x 5° grids, and all spatial data were available at this resolution.

Three sources of spatial data were used:

- 1) The ocean area of each grid  $g$  in each region  $r$ :  $A_{rg}$  accounting for area variation with latitude, and non-ocean (land) area. This was calculated using the R code in Appendix A. The projection was the Lambert azimuthal equal-area projection.
- 2) The year-invariant relative density of porbeagle shark by grid and quarter  $q$  so that  $\sum_g D_{rgq} = 1$  for all quarters across regions (see Section 2.3). Because no density estimates were available south of 45 °S, we assumed that the density at each longitude was constant south of the most southerly estimate, with the distribution limited at 55 °S (Figure 18). The true spatial distribution is uncertain.
- 3) The absolute fishing effort between 1960 and 2015, by grid and quarter, summed over all fleets combined (Figure 18). All effort was converted to the number of sets, with each set standardised to 3000 hooks. This is the assumed effort unit for the Japanese CPUE data.




Figure 18. Illustrative map of five regions showing (top) the spatial coverage of relative density data of porbeagle shark, averaged across quarters, and (bottom) the sum of absolute fishing effort between 1960 and 2015. Scale is from red (high) to yellow (low).

### 2.6.2 Fitting of biomass dynamic model to catch and abundance in the calibration area

Relative abundance indices from the Western Indian/Eastern Atlantic Ocean region (section 2.5.1) were used to calibrate a population catchability parameter for porbeagle shark over the spatial domain of the assessment. To estimate a posterior distribution for the catchability coefficient  $q$ , we fitted a biomass dynamic model (BDM) to Japanese observer CPUE index for the calibration region (Western Indian/Eastern Atlantic Ocean region), using a reconstructed catch for the whole region (Section 2.4, Figure 12).

The biomass dynamic, state-space model was implemented using the R package ‘bdm’ (Edwards 2017) which is written in the Bayesian modelling language Stan (Stan Development Team 2014).

The model takes the form:

$$\begin{aligned}\mu_{t+1} &= x_t + r \cdot x_t \cdot (1 - x_t) - C_t/K - \sigma_{[p]}^2/2 \\ x_{t+1} &\sim LN(\mu_{t+1}, \sigma_{[p]}^2) \\ I_t &\sim LN(x_t, \sigma_{[o]}^2)\end{aligned}$$

where  $x$  is the unobserved biomass depletion relative to the carrying capacity  $K$ ,  $C$  is the catch,  $\mu$  is the expected value (i.e.  $\mu = E[x]$ ),  $I$  is the observed CPUE index,  $\sigma_{[p]}^2$  is process variance,  $\sigma_{[o]}^2$  is observation variance, and  $q$  is the catchability scalar.  $\sim LN$  indicates a lognormal distribution. The term ‘ $-\sigma_{[p]}^2/2$ ’ is an adjustment for lognormal bias. We assumed a uniform prior on  $\log(K)$  with a plausible upper bound (based on expert opinion) equivalent to 5 individuals per km<sup>2</sup>. This was a relatively arbitrary value designed to be consistent with the  $K$  upper limit of 1 per km<sup>2</sup> assumed for the much rarer bigeye thresher (Fu et al. 2016). The prior for the intrinsic growth rate  $r$  was derived from life-history data (Table 2) (prior mean=0.033, cv=0.55). The model was run with 4 chains of 2000 samples each. A burn-in period of 1000 samples from each chain was discarded, leaving 4000 samples in total.

Table 2. Input life history information used to develop a prior for the maximum intrinsic population growth rate ( $r$ ) of porbeagle shark within the spatial domain of the assessment. Maturation, growth and recruitment parameters were based on available information for females only. Parameter values were reviewed by Clarke et al. (2015) and those values were either adopted here (with original sources given) or modified according to the listed source.

| Process       | Parameter               | Value       | CV   | Reference                                                                                                                                                                                                                  |
|---------------|-------------------------|-------------|------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Longevity     | $A_{inf}$ (yr)          | 75          |      | Francis et al. (2007)                                                                                                                                                                                                      |
| Maturation    | $A_{50}$ (yr)           | 14.5        | 0.25 | Francis (2015)                                                                                                                                                                                                             |
|               | delta                   | 1.5         | 0.25 |                                                                                                                                                                                                                            |
| Growth        | $L_{inf}$ (cm, FL)      | 211         | 0.3  | Francis (2015)                                                                                                                                                                                                             |
|               | $k$                     | 0.086       | 0.3  | Francis (2015)                                                                                                                                                                                                             |
|               | $t_0$                   | -6.1        | 0.3  | Francis (2015)                                                                                                                                                                                                             |
|               | $a$                     | 2.14289E-05 | 0.1  | Ayers et al. (2004)                                                                                                                                                                                                        |
| Length-weight | $b$                     | 2.924       | 0.1  |                                                                                                                                                                                                                            |
| Recruitment   | $\alpha$ (no.)          | 3.75        |      | Clarke et al. (2015)                                                                                                                                                                                                       |
| Mortality     | $M$ (yr <sup>-1</sup> ) | 0.09        | 0.42 | Averaged from four empirical equations: 1. Hoenig (1983): $\ln(M)=0.941-0.873 \ln(A_{max})$ ; 2. Campana et al. (2001): $M=-\ln(0.01/A_{max})$ ; 3. Jensen (1996): $M=1.65/Amat$ ( $=A_{50}$ ); 4. Jensen (1996): $M=1.6k$ |

The catch and abundance data are shown in Figure 19. Trace outputs from the Monte Carlo Markov Chain (MCMC) model fit, the derived fit to the CPUE index, and the posterior distributions of  $q$ ,  $r$  and  $\log(K)$  are shown in Figure 20.




Figure 19. Catch and abundance data.




Figure 20. Top: Trace outputs from the MCMC model fit showing estimates of  $r$  and  $\log(K)$  alongside the log-posterior ( $lp\_$ ). Middle: Derived fit to the CPUE index. Bottom: Posterior distributions of  $q$ ,  $r$  and  $\log(K)$ .

### 2.6.3 Estimation of the fishing mortality for each assessment region

We first convert the catchability estimated using the BDM into a catchability that can be used in the risk assessment. For BDM the catch equation is:

$$C = q^{[bdm]} \cdot \frac{N}{K} \cdot E$$

where  $N$  is the total number in the assessment area,  $K$  is the carrying capacity (total number), and  $E$  is the effort in set units; whereas for the fisheries risk assessment ( $fra$ ), the catch is a function of the relative density in numbers per unit area  $D = N/A$ , with area expressed in  $\text{km}^2$ , rather than the depletion  $N/K$ , and the catch equation is therefore:

$$C = q^{[fra]} \cdot \frac{N}{A} \cdot E$$

Our estimate of the catchability for the risk assessment is therefore:

$$q^{[fra]} = q^{[bdm]} \cdot \frac{A}{K} = q^{[bdm]} \cdot \frac{1}{\tilde{K}}$$

where  $\tilde{K}$  is the carrying capacity expressed in number per  $\text{km}^2$  (i.e. the absolute density at carrying capacity). The area  $A$  is taken to be a summation across grids with non-zero density (i.e.  $A = \sum_g A_g |D_g > 0$ ). We ignore seasonality in our calculation of  $A$  because any grid with non-zero density is non-zero for the whole year. This derivation of the catchability is valid for all of the assessment regions, using the approximation that the density at carrying capacity is constant across regions.

To derive a fishing mortality, we first estimate catches by year  $y$ , grid  $g$ , quarter  $q$  and iteration  $i$  for each posterior sample of the catchability, where catch  $C$  is a function of the relative density  $D$  and effort  $E$ :

$$C_{yqqi} = q_i^{[fra]} \cdot D_{gq} \cdot E_{yqq}$$

We then estimate total numbers for each year, grid and quarter as the product of the relative density and the area  $A_g$  in  $\text{km}^2$  for each grid:

$$N_{gq} = D_{gq} \cdot A_g$$

To estimate the harvest rate  $U$  using the risk assessment methodology, we calculate:

$$U_{yqi} = \frac{\sum_g C_{yqqi}}{\sum_g N_{gq}}$$

Assuming an exponential model of instantaneous mortality we can then write the fishing mortality as:

$$F_{yqi} = -\ln(1 - U_{yqi})$$

Finally, we calculate the fishing mortality rate (impact) per year by taking the sum across quarters:

$$F_{yi} = \sum_q F_{yqi}$$

Annual fishing mortalities are shown in Figure 21 and Appendix B. Annual  $F$  values were greatest in Western Indian/Eastern Atlantic Ocean, slightly lower in the Eastern Indian Ocean, and lowest in the



Western Pacific Ocean. Annual median  $F$  decreased from the mid-1980s to 2014 in both the Western Indian Ocean/Eastern Atlantic Ocean and Eastern Indian Ocean regions. In the assessment area (three regions combined) in the last decade (2005 to 2014), median  $F$  values ranged from 0.0008 to 0.0015 (mean 0.0010). Higher  $F$  values in the Western Indian Ocean/Eastern Atlantic Ocean and Western Pacific Ocean regions were consistently associated with the second quarter (April-June months or austral autumn season) (quarterly  $F$  estimates not shown). In contrast, the third and fourth quarters (July-September (austral winter) and October-December (austral spring)) contributed higher  $F$  values in the Eastern Indian Ocean region, however with a shift to higher  $F$  values in the second and third quarter (and a drop in fourth quarter  $F$ ) over the recent period (2010-2014).

#### 2.6.4 Estimation of risk

Risk  $R$  is calculated as the ratio between impact (sum of spatially-explicit relative  $F$  estimates) and our limit reference point  $F^{[CRASH]}$  for the stock (the MIST). This is the fishing mortality that will lead to a population crash, and was set equal to the posterior distribution of  $r$  as obtained from the BDM fit. The risk is then:

$$R = \frac{Impact}{MIST} = \frac{F}{F^{[CRASH]}}$$

which was calculated for each year using the full posterior distributions of both  $F$  and  $F^{[CRASH]}$ .

Risk values are shown both as an  $F$ -ratio and the probability that  $F$  exceeds the MIST in Figures 22 and 23, and for the period from 1992 onwards (the first year of Japanese CPUE data) in Tables 3 and 4.  $F$ -ratios for the assessment area declined by half from a mean of 0.068 (range 0.051–0.088) in 1992-2005, to a mean of 0.032 (range 0.023–0.042) in 2006-2014 (Table 3). The probability of  $F$  exceeding the MIST decreased by 95% from a mean of 0.0084 (range 0.0015–0.0205) in 1992-2005, to a mean of 0.0004 (range 0.0000–0.0013) in 2006-2014 (Table 4).

#### 2.6.5 Contributions to fishing mortality

We estimated the proportional contributions to fishing mortality of each fishery by year in each region, and for the different fishing strategies within the pelagic longline fishery. Fishing strategies (which we have also called ‘fisheries’) were determined based on cluster analysis of species composition data in the Japanese fleet (see Hoyle et al. (2017) for a description of the approach). For each grid cell by year-quarter stratum, the Japanese longliners’ proportional allocations of effort by cluster were noted, and assigned to all effort in that stratum. For effort records in strata that included no Japanese effort, the same approach was applied at coarser stratification: latitude (within region) by year-quarter. For the few records still not assigned, the same approach was applied with stratification at latitude (within region) by quarter.

Predicted longline catch was then calculated by region, year, and cluster by summing across grid cells and quarters. Other catch types available as total catch were then added to the dataset. Proportional contributions to  $F$  were calculated as  $\text{catch}(\text{fishery, year}) / \text{total catch}(\text{year})$ . Please note that uncertainties in predicted catches can significantly affect these estimates of proportional contributions to  $F$ , which should be regarded as approximate and indicative.

The greatest contributions to  $F$  were made by the pelagic longline fisheries, with the largest contribution by the SBT fishery and the mixed ALB/SBT fisheries (Figure 24). The contribution of ALB/SBT increased significantly from about 2004 in the Western Indian / Eastern Atlantic Ocean region, and from 2005 in the Eastern Indian Ocean region, but has been consistently important in the

Western Pacific since 1996. The more northern longline fishery that takes albacore, bigeye and yellowfin tunas made a generally smaller but still substantial contribution in recent years. In the Western Pacific region, the New Zealand midwater trawl fishery has also contributed to the fishing mortality.




Figure 21. Estimated fishing mortalities by year and region. The horizontal dotted line indicates the median MIST value, but note that the MIST varies among realisations of the model, with the grey band showing the 90% distribution. The boxes show the interquartile range of the  $F$  estimates, with a line at the median of each. The whiskers extend up to 1.5 x the interquartile range. Dots mark points beyond the whiskers.




Figure 22:  $F$ -ratio plots showing the median values of  $F / MIST$  by year, for three regions, and for the three regions combined (the assessment area). Note that the  $F$ -ratio is always below 1, which is the upper bound of the  $y$  axis.




Figure 23: Risk plots showing the probability that  $F$  exceeds the MIST by year, for three regions, and for the three regions combined (the assessment area).

Table 3: *F*-ratio metric for the impact of pelagic longline fisheries on porbeagle shark in three regions, and in the combined regions (the assessment area). The *F*-ratio metric is the ratio of *F* to the limit reference point (MIST), rounded to three decimal places. 95% confidence intervals for these values are shown in Appendix C.

| Year | Western Indian Ocean/Eastern Atlantic Ocean | Eastern Indian Ocean | Western Pacific Ocean | Combined |
|------|---------------------------------------------|----------------------|-----------------------|----------|
| 1992 | 0.088                                       | 0.090                | 0.067                 | 0.088    |
| 1993 | 0.132                                       | 0.036                | 0.053                 | 0.080    |
| 1994 | 0.095                                       | 0.045                | 0.037                 | 0.061    |
| 1995 | 0.090                                       | 0.055                | 0.032                 | 0.063    |
| 1996 | 0.074                                       | 0.073                | 0.024                 | 0.064    |
| 1997 | 0.082                                       | 0.077                | 0.033                 | 0.073    |
| 1998 | 0.081                                       | 0.086                | 0.049                 | 0.075    |
| 1999 | 0.094                                       | 0.078                | 0.054                 | 0.079    |
| 2000 | 0.065                                       | 0.092                | 0.041                 | 0.073    |
| 2001 | 0.095                                       | 0.074                | 0.061                 | 0.081    |
| 2002 | 0.063                                       | 0.043                | 0.065                 | 0.053    |
| 2003 | 0.050                                       | 0.055                | 0.056                 | 0.051    |
| 2004 | 0.071                                       | 0.051                | 0.031                 | 0.055    |
| 2005 | 0.070                                       | 0.043                | 0.021                 | 0.051    |
| 2006 | 0.062                                       | 0.030                | 0.011                 | 0.037    |
| 2007 | 0.045                                       | 0.031                | 0.012                 | 0.031    |
| 2008 | 0.046                                       | 0.039                | 0.011                 | 0.038    |
| 2009 | 0.052                                       | 0.044                | 0.013                 | 0.042    |
| 2010 | 0.037                                       | 0.035                | 0.011                 | 0.030    |
| 2011 | 0.042                                       | 0.029                | 0.013                 | 0.030    |
| 2012 | 0.033                                       | 0.019                | 0.015                 | 0.023    |
| 2013 | 0.032                                       | 0.029                | 0.017                 | 0.028    |
| 2014 | 0.022                                       | 0.038                | 0.017                 | 0.026    |

Table 4: Risk metric for the impact of pelagic longline fisheries on porbeagle shark in three regions, and in the combined regions (the assessment area). The risk metric is the probability that  $F$  is greater than the limit reference point (MIST), rounded to four decimal places. 95% confidence intervals for these values are shown in Appendix C.

| Year | Western Indian Ocean/Eastern Atlantic Ocean | Eastern Indian Ocean | Western Pacific Ocean | Combined |
|------|---------------------------------------------|----------------------|-----------------------|----------|
| 1992 | 0.0170                                      | 0.0190               | 0.0060                | 0.0205   |
| 1993 | 0.0450                                      | 0.0013               | 0.0038                | 0.0115   |
| 1994 | 0.0195                                      | 0.0018               | 0.0008                | 0.0065   |
| 1995 | 0.0198                                      | 0.0043               | 0.0003                | 0.0063   |
| 1996 | 0.0110                                      | 0.0100               | 0.0000                | 0.0070   |
| 1997 | 0.0130                                      | 0.0115               | 0.0008                | 0.0090   |
| 1998 | 0.0130                                      | 0.0160               | 0.0020                | 0.0095   |
| 1999 | 0.0208                                      | 0.0108               | 0.0033                | 0.0118   |
| 2000 | 0.0088                                      | 0.0218               | 0.0015                | 0.0090   |
| 2001 | 0.0248                                      | 0.0120               | 0.0075                | 0.0148   |
| 2002 | 0.0055                                      | 0.0010               | 0.0075                | 0.0018   |
| 2003 | 0.0025                                      | 0.0045               | 0.0035                | 0.0035   |
| 2004 | 0.0090                                      | 0.0035               | 0.0003                | 0.0045   |
| 2005 | 0.0105                                      | 0.0018               | 0.0000                | 0.0015   |
| 2006 | 0.0053                                      | 0.0008               | 0.0000                | 0.0008   |
| 2007 | 0.0025                                      | 0.0003               | 0.0003                | 0.0003   |
| 2008 | 0.0010                                      | 0.0008               | 0.0000                | 0.0008   |
| 2009 | 0.0033                                      | 0.0025               | 0.0000                | 0.0013   |
| 2010 | 0.0005                                      | 0.0008               | 0.0000                | 0.0000   |
| 2011 | 0.0013                                      | 0.0003               | 0.0000                | 0.0000   |
| 2012 | 0.0000                                      | 0.0000               | 0.0000                | 0.0003   |
| 2013 | 0.0003                                      | 0.0000               | 0.0000                | 0.0003   |
| 2014 | 0.0000                                      | 0.0013               | 0.0000                | 0.0000   |




Figure 24: Proportional contributions by year and region to fishing mortality, for the fisheries operating in each region. The pelagic longline effort is broken into three fishing strategies based on cluster analysis (see Hoyle et al. 2017). Note that the French (Kerguelen and Crozet Islands) and Australian (Heard Island and McDonald Islands) sub-Antarctic fisheries' contributions to  $F$  are too small to be visible on the plots.

## 2.7 Quantitative stock assessment

We explored the potential to develop a quantitative stock assessment model using Stock Synthesis (Methot & Wetzel 2013) for the porbeagle shark stock. Age structured approaches have the advantage that they may provide more reliable inferences than simpler biomass dynamic models by using more data types and modelling important processes that biomass dynamic models cannot represent. For example, when selectivity changes through time it changes the productivity of the population. Representing such changes is difficult with a biomass dynamic model, and failing to represent them often introduces bias (Wang et al. 2014). However, age structured models have the disadvantages that they require more data than biomass dynamic models, and are time-consuming to develop.

The porbeagle shark stock has significant spatial size structure, with smaller sharks observed in warmer waters in the north and larger sharks further south. Furthermore, the northern areas with smaller sharks experience more fishing effort than the rest of the distribution, so selectivity is expected to be biased towards smaller sharks. There is potential to estimate selectivity parameters, because information on catch at size is available from observer data on size (and sex) in the Japanese, New Zealand, Argentinian, and Chilean fisheries. There is also information on relationships between capture size in longline fisheries and SST for the Japanese fishery (Hoyle et al. 2017b), which could be used to impute the size structure taken by all longline fisheries, which together dominate the catch.

On the other hand, the spatial distribution of fishing effort has been relatively stable through time, so it is likely that changes in selectivity through time have been relatively small. It is thus anticipated that this source of potential bias is not a major influence on the results of the assessment.

The decision to structure the spatial domain of the risk assessment model as multiple regions (rather than a single region), implies the need for a multi-regional age structured model. This additional work on the age structured model proved impractical in this study, however, it could be valuable in future to explore alternative hypotheses regarding fishery selectivity and population structure.

## 3 Discussion

This risk assessment for porbeagle shark in the Southern Hemisphere treats the population as five separate subpopulations, and provides estimates of the risk of exceeding the MIST limit reference point for the three subpopulations with sufficient data. Risk was also estimated for the three together. The remaining two subpopulations have less data, and their status has been interpreted based on relative abundance indices.

Results indicate low fishing mortality rates in the three regions comprising the assessment area, and low risk from commercial pelagic longline fisheries to porbeagle shark over the spatial domain of the assessment. These results are consistent with the trends observed in catch rate indicators over the entire Southern Hemisphere range of the porbeagle shark population, which in most cases show stable or increasing catch rates. Concern has previously been expressed about reduced catch rates in the Uruguay longline fishery after 1993 (International Commission for the Conservation of Atlantic Tunas 2010), but the re-analysis undertaken for this study in collaboration with Uruguayan researchers indicates that in 1993 both the vessels fishing and their fishing methods changed almost completely (Forselledo et al. 2017). After allowing for this change, a decline was no longer evident.



Most catch rate indicators were relatively short, variable, and uncertain, with the majority either stable or increasing. Length indicators were also variable. Only the Argentinian size and sex indicators showed temporal trends, with a small decline in sizes for both sexes, and a slight trend towards less female bias in the sex ratio index.

The indicator analyses, in addition to providing time series to monitor population change, revealed spatial patterns in size and sex distributions, and relationships with environmental variables. Such analyses are critical inputs to stock status assessments, because they help to determine model structure.

The catch rate indicators are by far the most important inputs to this status assessment, and their reliability determines the reliability of the assessment. Stable or increasing observed population trends, under fishing pressure, constrain the risk assessment model to estimate levels of catchability and population density that would allow the population to be stable or increasing. Thus the indicator trend in the calibration area is the most important factor determining the relatively low estimate of risk. Continued data collection by observers will improve the time series and provide better evidence about abundance trends. Maintaining collection and analysis of indicators from observer data is a key recommendation from this project.

Furthermore, the population catchability was calibrated assuming that capture mortality was 100% (i.e., zero post-release survival). In recent years many fleets have released porbeagles, and many of these released sharks are likely to have survived (Campana et al. 2016). Allowing for post-release survival would reduce these fishing mortality estimates, and reduce the estimated risk below the low risk levels estimated here.

The risk assessment assumes that catchability estimates from Japanese vessels are applicable to other fleets. The three targeting strategies identified for the Japanese fleet, which were explored by using clusters in the CPUE standardisation, had quite similar catch rates. This observation provides some reassurance about the applicability of our catch rate estimates to other fleets. Although it is possible that some other fleets' targeting strategies may have very different catch rates for porbeagles, we have no evidence for this.

This approach also assumes that SST estimates from observers on Japanese vessels are equivalent to modelled SST estimates from the CSIRO CARS database, the CARS estimates being averages across multiple years with no allowance for inter-year variability. There is likely to be significant divergence between local, vessel-based observations of SST, and averaged model predictions, but we do not think this sufficient to substantially change the stock status results.

We also assumed an upper prior bound for  $K$  equivalent to a maximum plausible density of 5 sharks per km<sup>2</sup>. The correct level for this assumption was difficult to determine precisely, and alternative values may slightly affect the outcomes.

The risk assessment results are based on strong assumptions about the population density distribution. We have assumed that density is driven primarily by SST, with smaller effects due to season, latitude and year, which implies that depletion is the same in all areas. This assumption was required to extend the analysis to areas and times without reliable catch data and thus cover the entire spatial domain. However, it is likely that spatial variation in historical effort has depleted some areas more, which would consequently have lower densities than other areas, and lower densities than we have assumed. Since catch is the product of effort and density, we would also have overestimated their pelagic longline catches. However, given the approach used in the assessment,

this effect does not bias the estimates of fishing mortality. Further exploration of this issue could involve applying our SST-based spatial-environmental model to the Japanese longline data by region, predicting distribution and catches independently for each region, and allowing re-estimation of catchability for the calibration region.

The risk assessment assumes that population density from 45 to 55 °S is the same as at 40 to 45 °S, since we have evidence from fisheries and surveys that they occur at these latitudes, but do not have Japanese longline observer data with which to estimate density. This is an important assumption, because it implies that the low fishing effort south of 45 °S provides a refuge from fishing mortality for the population. Unfortunately there is little information about patterns of population density south of 45 °S, but there is considerable evidence that porbeagles are found there, with observations recorded for most of the regions considered here. In the Eastern Pacific catch rates were observed increasing to the southern limits of the JAMARC longline survey (60 °S) and were relatively stable to the southern limits of the JAMARC gillnet survey (52.5 °S) (Yatsu 1995, Semba et al. 2013, Hoyle et al. 2017b). Also in the Eastern Pacific, porbeagles have been caught in the Chilean demersal longline fishery to at least 56 °S. In the Western Pacific they have been taken in the New Zealand midwater trawl fishery to 53 °S (Francis 2013). In the Western Indian Ocean, porbeagles have been taken in the mackerel icefish and Patagonian toothfish trawl and longline fisheries in the Heard and McDonald Island EEZ (53 °S) (ABARES data, Heather Chapman, pers. comm.). The analysis of data from the Western Atlantic Argentinian surimi fleet (Cortés et al. 2017) conducted for this study shows catch rates increasing to the south, but this result may be due to the local geography, with strong currents, temperature variation, and depth changes around Cape Horn. We recommend further work to understand this southerly population, such as future analysis of bycatch information being collected from the Chilean demersal longline fishery, which fishes as far south as 56 °S.

We also recommend exploring selectivity at age in the Japanese pelagic longline data, which may permit estimation of the availability at age of the population to fishing. This analysis may permit two further developments: an age-structured version of the bdm biomass dynamic risk assessment (Edwards 2017); and direct estimation of the proportion of the population south of 45 °S, removing the need to assume constant density from 45 to 55 °S.

This analysis assumed separation of the population into five regions, but there is little information available with which to determine appropriate stock boundaries. We recommend analyses of distribution using various types of markers (genetics, microchemistry, stable isotopes, parasites, conventional and electronic tags) to identify biologically-based boundaries.

The multiple indicators/risk assessment approach served to 1) source and synthesise available information on porbeagle shark at the scale of the Southern Hemisphere; 2) identify important data gaps (e.g., density distribution and life-stage specific vulnerability and overlap with fishing activities); 3) define a productivity-based reference point for the species; and 4) prioritise fishery areas for monitoring and management. This project has filled important information gaps by both directly analysing available life history information, and providing statistical support to the analyses by participating national fisheries scientists.

The project has provided the first assessment of the sustainability of the impact of fishing on the Southern Hemisphere porbeagle shark stock, and laid a foundation for future work. Results indicate that the impact of fishing is low across the entire Southern Hemisphere range of the porbeagle shark population.

## 4 Acknowledgements

This project was funded by the Common Oceans (Areas Beyond National Jurisdiction (ABNJ)) Tuna Project.

We thank our collaborators as follows:

Japan: National Research Institute of Far Seas Fisheries, Yasuko Semba, Mikihiro Kai, Hiroaki Okamoto and Yujiro Akatsuka

New Zealand: Ministry for Primary Industries and Kath Large

Argentina: INIDEP, Federico Cortés, Juan Waessle and Ana Massa

Uruguay: DINARA, Rodrigo Forselledo, Federico Mas and Andres Domingo

Chile: IFOP, Juan Carlos Quiroz, Patricia Zarate, Daniel Devia, Jorge Azocar and Carlos Bustamante

France: Guy Duhamel, Muséum National d'Histoire Naturelle

Australia: Heather Chapman, ABARES

SPRFMO: Craig Loveridge

ICCAT: Paul de Bruyn

SPC: Peter Williams

CCSBT: Bob Kennedy and Colin Millar

## 5 References

- Australian Fisheries Management Authority (2017). Sharks – how AFMA and industry minimise interactions. <<http://www.afma.gov.au/portfolio-item/sharks>> (Accessed on 3 July 2017).
- Ayers, D.; Francis, M.P.; Griggs, L.H.; Baird, S.J. (2004). Fish bycatch in New Zealand tuna longline fisheries, 2000-01 and 2001-02. *New Zealand fisheries assessment report No. 2004/46*. 47 p.
- Biais, G.; Coupeau, Y.; Seret, B.; Calmettes, B.; Lopez, R.; Hetherington, S.; Righton, D. (2017). Return migration patterns of porbeagle shark (*Lamna nasus*) in the Northeast Atlantic: implications for stock range and structure. *ICES journal of marine science* 74: 1268–1276.
- Braccini, J.M.; Gillanders, B.M.; Walker, T.I. (2006). Hierarchical approach to the assessment of fishing effects on non-target chondrichthyans: case study of *Squalus megalops* in southeastern Australia. *Canadian journal of fisheries and aquatic sciences* 63: 2456-2466.
- Bruce, B. (2014). Shark futures: a synthesis of available data on mako and porbeagle sharks in Australasian waters. Current status and future directions. *FRDC Project No. 2011/045*.
- Campana, S.E.; Gibson, A.J.F.; Fowler, M.; Dorey, A.; Joyce, W. (2010a). Population dynamics of porbeagle in the northwest Atlantic, with an assessment of status to 2009 and projections for recovery. *Collective volume of scientific papers ICCAT 65*: 2109-2182.
- Campana, S.E.; Gibson, A.J.F.; Fowler, M.; Dorey, A.; Joyce, W. (2012). Population dynamics of Northwest Atlantic porbeagle (*Lamna nasus*), with an assessment of status and projections for recovery. *Canadian Science Advisory Secretariat research document No. 2012/096*. 84 p.
- Campana, S.E.; Joyce, W.; Fowler, M. (2010b). Subtropical pupping ground for a cold-water shark. *Canadian journal of fisheries and aquatic sciences* 67: 769-773.
- Campana, S.E.; Joyce, W.; Fowler, M.; Showell, M. (2016). Discards, hooking, and post-release mortality of porbeagle (*Lamna nasus*), shortfin mako (*Isurus oxyrinchus*), and blue shark (*Prionace glauca*) in the Canadian pelagic longline fishery. *ICES journal of marine science* 73: 520–528.
- Campana, S.E.; Joyce, W.; Marks, L.; Hurley, P.; Natanson, L.J.; Kohler, N.E.; Jensen, C.F.; Mello, J.J.; Pratt, H.L.; Myklevoll, S.; Harley, S. (2008). The rise and fall (again) of the porbeagle shark population in the Northwest Atlantic. In: Camhi, M.D.; Pikitch, E.K.; Babcock, E.A. (eds). *Sharks of the open ocean: biology, fisheries and conservation*, pp. 445-461. Blackwell Publishing, Oxford, United Kingdom.
- Campana, S.E.; Marks, L.; Joyce, W.; Harley, S. (2001). Analytical assessment of the porbeagle shark (*Lamna nasus*) population in the Northwest Atlantic, with estimates of long-term sustainable yield. *Canadian Science Advisory Secretariat research document No. 2001/067*. 59 p.
- Clarke, S.; Coelho, R.; Francis, M.; Kai, M.; Kohin, S.; Liu, K.-M.; Simpfendorfer, C.; Tovar-Avila, J.; Rigby, C.; Smart, J. (2015). Report of the Pacific Shark Life History Expert Panel Workshop, 28-30 April 2015. *Western Central Pacific Fisheries Commission Scientific Committee eleventh regular session No. WCPFC-SC11-2015/EB-IP-13*. 111 p.
- Clarke, S.; Manarangi-Trott, L.; Brouwer, S. (2014). Issues for t-RFMOs in relation to the listing of shark and ray species by the Convention on International Trade in Endangered Species (CITES).

Western Central Pacific Fisheries Commission Scientific Committee tenth regular session No. WCPFC-SC10-2014/EP-IP-05. [http://www.wcpfc.int/system/files/EB-IP-05%20\(CITES%20%20Sharks\).pdf](http://www.wcpfc.int/system/files/EB-IP-05%20(CITES%20%20Sharks).pdf).

Clarke, S.C.; Francis, M.P.; Griggs, L.H. (2013). Review of shark meat markets, discard mortality and pelagic shark data availability, and a proposal for a shark indicator analysis. *New Zealand fisheries assessment report No. 2013/65*. 74 p.

Commission for the Conservation of Antarctic and Marine Living Resources (2006). Conservation Measure 32-18: Conservation of Sharks. <<https://www.ccamlr.org/en/measure-32-18-2006>> (Accessed on 3 July 2017).

Compagno, L.J.V. (2001). Sharks of the world. An annotated and illustrated catalogue of shark species known to date. *FAO species catalogue for fishery purposes No. 1, vol. 2*. 269 p.

Convention on Migratory Species (2016). Memorandum of Understanding on the Conservation of Migratory Sharks (Shark MOU). Fact Sheet. <[http://www.cms.int/sharks/sites/default/files/publication/fact\\_sheet\\_sharks\\_english\\_0.pdf](http://www.cms.int/sharks/sites/default/files/publication/fact_sheet_sharks_english_0.pdf)> (Accessed on 3 July 2017).

Cortés, E. (2002). Incorporating uncertainty into demographic modeling: application to shark populations and their conservation. *Conservation biology* 16: 1048-1062.

Cortés, E. (2008). Comparative life history and demography of pelagic sharks. *In*: Camhi, M.D.; Pikitch, E.K.; Babcock, E.A. (eds). *Sharks of the open ocean: biology, fisheries and conservation*, pp. 309-322. Blackwell Publishing, Oxford, United Kingdom.

Cortés, E.; Arocha, F.; Beerkircher, L.; Carvalho, F.; Domingo, A.; Heupel, M.; Holtzhausen, H.; Neves, M.; Ribera, M.; Simpfendorfer, C. (2010). Ecological risk assessment of pelagic sharks caught in Atlantic pelagic longline fisheries. *Aquatic living resources* 23: 25–34.

Cortés, E.; Domingo, A.; Miller, P.; Forselledo, R.; Mas, F.; Arocha, F.; Campana, S.; Coelho, R.; Da Silva, C.; Hazin, F.H.V.; Holtzhausen, H.; Keene, K.; Lucena, F.; Ramirez, K.; Santos, M.N.; Semba-Murakami, Y.; Yokawa, K. (2015). Expanded ecological risk assessment of pelagic sharks caught in Atlantic pelagic longline fisheries. *Collective volume of scientific papers ICCAT 71*: 2637-2688.

Cortés, F.; Waessle, J.A.; Massa, A.M.; Hoyle, S.D. (2017). Aspects of porbeagle shark bycatch in the Argentinean surimi fleet operating in the Southwestern Atlantic Ocean (50 – 57° S) during 2006-2014. *WCPFC Scientific Committee 13th regular session WCPFC-SC13-SA-IP-14*.

Curtis, T.H.; Laporte, S.; Cortes, E.; DuBeck, G.; McCandless, C. (2016). Status review report: porbeagle shark (*Lamna nasus*). Final report to National Marine Fisheries Service, Office of Protected Resources. February 2016. <[http://www.greateratlantic.fisheries.noaa.gov/protected/pcp/soc/porbeagle\\_shark.html](http://www.greateratlantic.fisheries.noaa.gov/protected/pcp/soc/porbeagle_shark.html)> (Accessed on 3 July 2017).

Ebert, D.A.; Fowler, S.; Compagno, L. (2013). *Sharks of the world. A fully illustrated guide*. Wild Nature Press, Plymouth, England. 528 p.

Edwards, C. (2017). bdm: Bayesian biomass dynamics model. R package version 0.0.0.9019. p.

European Union (2015). Council Regulation (EU) 2015/104 of 19 January 2015 fixing for 2015 the fishing opportunities for certain fish stocks and groups of fish stocks, applicable in Union waters and, for Union vessels, in certain non-Union waters, amending Regulation (EU) No 43/2014 and repealing Regulation (EU) No 779/2014. <<http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A32015R0104>> (Accessed on 3 July 2017).

Food and Agriculture Organization of the United Nations (2012). Recommendation GFCM/36/2012/3 on fisheries management measures for conservation of sharks and rays in the GFCM area. <<http://www.fao.org/3/a-ax385e.pdf>> (Accessed on 3 July 2017).

Food and Agriculture Organization of the United Nations (2017). Aquatic species fact sheet: *Lamna nasus* (Bonnaterre, 1788). <<http://www.fao.org/fishery/species/search/en>> (Accessed on 3 July 2017).

Forselledo, R.; Mas, F.; Hoyle, S.D.; Domingo, A. (2017). Standardized cpue of porbeagle shark (*Lamna nasus*) caught by the Uruguayan pelagic longline fleet (1982-2012). *WCPFC Scientific Committee 13th regular session WCPFC-SC13-SA-IP-18*.

Francis, M.; Large, K. (2017). Updated abundance indicators for New Zealand blue, porbeagle and shortfin mako sharks. *CCSBT Ecologically Related Species Working Group 12th meeting CCSBT-ERS/1703/14. WCPFC Scientific Committee 13th regular session WCPFC-SC13-2017/SA-IP-13*. 19 p.

Francis, M.P. (2013). Commercial catch composition of highly migratory elasmobranchs. *New Zealand fisheries assessment report No. 2013/68*. 79 p.

Francis, M.P. (2015). Size, maturity and age composition of porbeagle sharks observed in New Zealand tuna longline fisheries. *New Zealand fisheries assessment report No. 2015/16*. 30 p.

Francis, M.P. (2017). Recalculation of historical landings of porbeagle shark. *New Zealand fisheries assessment report 2017/12. WCPFC Scientific Committee 13th regular session WCPFC-SC13-2017/SA-IP-16*. 20 p.

Francis, M.P.; Campana, S.E.; Jones, C.M. (2007). Age under-estimation in New Zealand porbeagle sharks (*Lamna nasus*): is there an upper limit to ages that can be determined from shark vertebrae? *Marine and freshwater research* 58: 10-23.

Francis, M.P.; Clarke, S.C.; Griggs, L.H.; Hoyle, S.D. (2014). Indicator based analysis of the status of New Zealand blue, mako and porbeagle sharks. *New Zealand fisheries assessment report No. 2014/69*. 109 p.

Francis, M.P.; Duffy, C. (2005). Length at maturity in three pelagic sharks (*Lamna nasus*, *Isurus oxyrinchus*, and *Prionace glauca*) from New Zealand. *Fishery bulletin* 103: 489-500.

Francis, M.P.; Holdsworth, J.C.; Block, B.A. (2015). Life in the open ocean: seasonal migration and diel diving behaviour of Southern Hemisphere porbeagle sharks (*Lamna nasus*). *Marine biology* 162: 2305-2323.

Francis, M.P.; Natanson, L.J.; Campana, S.E. (2008). The biology and ecology of the porbeagle shark, *Lamna nasus*. In: Camhi, M.D.; Pikitch, E.K.; Babcock, E.A. (eds). *Sharks of the open ocean: biology, fisheries and conservation*, pp. 105-113. Blackwell Publishing, Oxford, United Kingdom.

- Francis, M.P.; Stevens, J.D. (2000). Reproduction, embryonic development and growth of the porbeagle shark, *Lamna nasus*, in the south-west Pacific Ocean. *Fishery bulletin* 98: 41-63.
- Fu, D.; Roux, M.-J.; Clarke, S.; Francis, M.; Dunn, A.; Hoyle, S. (2016). Pacific-wide sustainability risk assessment of bigeye thresher shark (*Alopias superciliosus*). *NIWA client report 2016089WN. WCPFC Scientific Committee 13th regular session WCPFC-SC13-2017/SA-WP-11*. 98 p.
- Gallagher, A.J.; Kyne, P.M.; Hammerschlag, N. (2012). Ecological risk assessment and its application to elasmobranch conservation and management. *Journal of fish biology* 80: 1727-1748.
- Griffiths, S.P.; Brewer, D.T.; Heales, D.S.; Milton, D.A.; Stobutzki, I.C. (2006). Validating ecological risk assessments for fisheries: assessing the impacts of turtle excluder devices on elasmobranch by-catch populations in an Australian trawl fishery. *Marine and freshwater research* 57: 395–401.
- Griggs, L.H.; Baird, S.J.; Francis, M.P. (2007). Fish bycatch in New Zealand tuna longline fisheries 2002-03 to 2004-05. *New Zealand fisheries assessment report No. 2007/18*. 58 p.
- Hoenig, J.M. (1983). Empirical use of longevity data to estimate mortality rates. *Fishery bulletin* 81: 898-903.
- Horn, P.L.; Ballara, S.L.; Sutton, P.J.H.; Griggs, L.H. (2013). Evaluation of the diets of highly migratory species in New Zealand waters. *New Zealand aquatic environment and biodiversity report No. 116*. 140 p.
- Hoyle, S.D.; Quiroz, J.C.; Zarate, P.; Devia, D.; Azocar, J. (2017a). Population indicators for porbeagle sharks in the Chilean swordfish fishery. *WCPFC Scientific Committee 13th regular session WCPFC-SC13-SA-IP-17*.
- Hoyle, S.D.; Semba, Y.; Kai, M.; Okamoto, H. (2017b). Development of Southern Hemisphere porbeagle shark stock abundance indicators using Japanese commercial and survey data. *New Zealand Fisheries Assessment Report 2017/07. WCPFC Scientific Committee 13th regular session WCPFC-SC13- SA-IP-15*. 64 p.
- International Commission for the Conservation of Atlantic Tunas. (2010). Report of the 2009 porbeagle stock assessments meeting. Copenhagen, Denmark, June 22 to 27, 2009. SCRS/2009/014. *Collective volume of scientific papers ICCAT 65*: 1909-2005.
- International Commission for the Conservation of Atlantic Tunas (2015). Recommendation by ICCAT on porbeagle caught in association with ICCAT fisheries. Recommendation 15-06. <https://www.iccat.int/Documents/Recs/compendiopdf-e/2015-06-e.pdf> (Accessed on 3 July 2017).
- Jensen, A.L. (1996). Beverton and Holt life history invariants result from optimal trade-off of reproduction and survival. *Canadian journal of fisheries and aquatic sciences* 53: 820-822.
- Kitamura, T.; Matsunaga, H. (2010). Population structure of porbeagle (*Lamna nasus*) in the Atlantic Ocean as inferred from mitochondrial DNA control region sequences. *Collective volume of scientific papers ICCAT 65*: 2082-2087.
- Last, P.R.; Stevens, J.D. (2009). Sharks and rays of Australia. Second. CSIRO, Hobart. 644 p.

Methot, R.D.; Wetzel, C.R. (2013). Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research* 142: 86-99.  
<<http://dx.doi.org/10.1016/j.fishres.2012.10.012>>

Ministry for Primary Industries. (2016). Fisheries Assessment Plenary: stock assessments and stock status, November 2016. Compiled by the Fisheries Science Group, Ministry for Primary Industries, Wellington, New Zealand. 459 p.

Murua, H.; Coelho, R.; Santos, M.N.; Arrizabalaga, H.; Yokawa, K.; Romanov, E.; Zhu, J.F.; Kim, Z.G.; Bach, P.; Chavance, P.; Delgado de Molina, A.; Ruiz, J. (2012). Preliminary ecological risk assessment (ERA) for shark species caught in fisheries managed by the Indian Ocean Tuna Commission (IOTC). *No. IOTC-2012-SC15-INF10 Rev\_1 (Dec 2012)*.

National Oceanic and Atmospheric Administration (2016). Notice of 12-month finding on petitions to list porbeagle shark as threatened or endangered under the Endangered Species Act (ESA). Federal Register Vol. 81, No. 147, 1 August 2016.  
<<http://www.greateratlantic.fisheries.noaa.gov/regs/2016/July/16porbeaglesharklistesa.pdf>>  
(Accessed on 3 July 2017).

Northeast Atlantic Fisheries Commission (2016). Recommendation on conservation and management measures for porbeagle (*Lamna nasus*) in the NEAFC Regulatory Area from 2016 to 2019. Recommendation 7:2016. <[https://www.neafc.org/system/files/Rec7\\_Porbeagle-from-2016-2019.pdf](https://www.neafc.org/system/files/Rec7_Porbeagle-from-2016-2019.pdf)> (Accessed on 3 July 2017).

Pade, N.G.; Queiroz, N.; Humphries, N.E.; Witt, M.J.; Jones, C.S.; Noble, L.R.; Sims, D.W. (2009). First results from satellite-linked archival tagging of porbeagle shark, *Lamna nasus*: area fidelity, wider-scale movements and plasticity in diel depth changes. *Journal of experimental marine biology and ecology* 370: 64–74.

Pons, M.; Domingo, A. (2010). Standardized CPUE of Porbeagle shark (*Lamna nasus*) caught by the Uruguayan pelagic longline fleet (1982-2008). *Collect. Vol. Sci. Pap. ICCAT No. 6*. 2098-2108 p.

R Core Team (2017). R: A language and environment for statistical computing [Internet]. Vienna, Austria; 2014. p.

Rice, J.; Tremblay-Boyer, L.; Scott, R.; Hare, S.; Tidd, A. (2015). Analysis of stock status and related indicators for key shark species of the Western Central Pacific Fisheries Commission. *Western Central Pacific Fisheries Commission Scientific Committee eleventh regular session No. WCPFC-SC11-2015/EB-WP-04-Rev 1*. <https://www.wcpfc.int/system/files/EB-WP-04%20shark%20indicators%20Rev%201.pdf>.

Ridgway, K.; Dunn, J.; Wilkin, J. (2002). Ocean interpolation by four-dimensional weighted least squares—Application to the waters around Australasia. *Journal of atmospheric and oceanic technology* 19(9): 1357-1375.

Saunders, R.A.; Royer, F.; Clarke, M.W. (2011). Winter migration and diving behaviour of porbeagle shark, *Lamna nasus*, in the Northeast Atlantic. *ICES journal of marine science* 68: 166-174.

Semba, Y.; Yokawa, K.; Matsunaga, H.; Shono, H. (2013). Distribution and trend in abundance of the porbeagle (*Lamna nasus*) in the southern hemisphere. *Marine and freshwater research* 64: 518-529.



Stan Development Team (2014). Stan: A C++ library for probability and sampling. Online: <http://mc-stan.org>. p.

Stevens, J.; Fowler, S.L.; Soldo, A.; McCord, M.; Baum, J.; Acuña, E.; Domingo, A.; Francis, M. (2006). *Lamna nasus*. The IUCN Red List of Threatened Species 2006: e.T11200A3261697. <<http://dx.doi.org/10.2305/IUCN.UK.2006.RLTS.T11200A3261697.en>> (Accessed on 3 July 2017).

Stobutzki, I.C.; Miller, M.J.; Heales, D.S.; Brewer, D.T. (2002). Sustainability of elasmobranchs caught as bycatch in a tropical prawn (shrimp) trawl fishery. *Fishery bulletin* 100: 800-821.

Testerman, C.; Richards, V.; Francis, M.; Pade, N.; Jones, C.; Noble, L.; Shivji, M. (2007). "Global phylogeography of the porbeagle shark (*Lamna nasus*) reveals strong genetic separation of Northern and Southern Hemisphere populations (Abstract)." Presented at the American Elasmobranch Society annual meeting, St Louis, USA.

Wang, S.-P.; Maunder, M.N.; Aires-da-Silva, A. (2014). Selectivity's distortion of the production function and its influence on management advice from surplus production models. *Fisheries Research* 158: 181-193.

Western and Central Pacific Fisheries Commission (2007). Rules and procedures for the protection, access to, and dissemination of data compiled by the Commission. *Western Central Pacific Fisheries Commission Scientific Committee Fourth Regular Session No. WCPFC4-2007-12*. 15 p.

Western and Central Pacific Fisheries Commission (2012). Process for designating WCPFC key shark species for data provision and assessment. <<https://www.wcpfc.int/system/files/Key-Doc-SC-08-Process-Designation-Key-WCPFC-Shark-Species.pdf>> (Accessed on 3 July 2017).

Williams, P.; Terawasi, P. (2016). Overview of tuna fisheries in the Western and Central Pacific Ocean, including economic conditions-2015. *Western Central Pacific Fisheries Commission Scientific Committee twelfth regular session No. WCPFC-SC12-2016/GN-WP-1 rev3*. <https://www.wcpfc.int/system/files/GN-WP-01%20Overview%20of%20WCPFC%20Fisheries%20Rev%203%20%286%20September%202016%29.pdf>

Wood, S.N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 73(1): 3-36.

Yatsu, A. (1995). Zoogeography of the epipelagic fishes in the South Pacific Ocean and the Pacific sector of the Subantarctic, with special reference to the ecological role of slender tuna, *Allothunnus fallai*. *Bulletin of the National Research Institute of Far Seas Fisheries* No. 32. 145 p.

Zhou, S.; Griffiths, S.P. (2008). Sustainability assessment for fishing effects (SAFE): a new quantitative ecological risk assessment method and its application to elasmobranch bycatch in an Australian trawl fishery. *Fisheries research* 91: 56-68.

## Appendix A R code for calculating areas of grid cells

```

windows()
m <- map(regions = c("South Africa", "Australia", "New Zealand", "Argentina",
"Chile", "Antarctica", "Uruguay", "Botswana", "Namibia", "Zimbabwe",
"South Georgia", "Falkland Islands", "Mozambique",
"Madagascar", "Brazil", "Paraguay", "French Southern and Antarctic Lands", "Heard
Island"), fill = TRUE)
#m <- map(fill = TRUE)
#identify.map(m)
res <- expand.grid(ln = seq(-180, 180, 5), lt = seq(-60, -30, 5), garea = NA, areax
= NA)
i=1
for (i in 1:length(res$lt)) {
  lt = res$lt[i]; ln = res$ln[i]
  geo_str <- paste0("+proj=laea +lon_0=",ln," +lat_0=",lt," +datum=WGS84")
  crs.geox <- CRS(geo_str)
  grcl <- data.frame(lon = c(ln, ln+5, ln+5, ln), lat = c(lt+5, lt+5, lt, lt))
  coordinates(grcl) <- ~ lon + lat
  projection(grcl) <- "+init=epsg:4326"
  grclx <- Polygon(grcl)

  cells <- Polygons(list(grclx), "onecell")
  cells2 <- SpatialPolygons(list(cells))
  projection(cells2) <- crs.geox

  m.spx <- map2SpatialPolygons(m, IDs=m$names,proj4string=crs.geox)
  m.spx <- gSimplify(m.spx, tol = 0.00001)
  m.spx <- gBuffer(m.spx, byid=TRUE, width=0)
  m.sp <- map2SpatialPolygons(m, IDs=m$names,proj4string=crs.geo)
  m.sp <- gSimplify(m.sp, tol = 0.00001)

  m.diff <- gDifference(cells2, m.sp)
  if(is.null(m.diff)) {
    res$garea[i] <- 0
    res$areax[i] <- 0
  } else {
    md2 <- spTransform(m.diff, crs.geox)
    res$garea[i] <- gArea(md2)/1e6
    res$areax[i] <- areaPolygon(m.diff)/1e6
  }
  print(i); flush.console()
}

```

## Appendix B      Fishing mortality estimates

Median  $F$  in each region and the assessment area (three regions combined) (rounded to four decimal places).

| Year | Western<br>Indian<br>Ocean/Eastern<br>Atlantic Ocean | Eastern<br>Indian<br>Ocean | Western<br>Pacific<br>Ocean | Combined |
|------|------------------------------------------------------|----------------------------|-----------------------------|----------|
| 1992 | 0.0026                                               | 0.0028                     | 0.0020                      | 0.0025   |
| 1993 | 0.0040                                               | 0.0011                     | 0.0016                      | 0.0024   |
| 1994 | 0.0028                                               | 0.0014                     | 0.0011                      | 0.0019   |
| 1995 | 0.0028                                               | 0.0016                     | 0.0009                      | 0.0019   |
| 1996 | 0.0022                                               | 0.0022                     | 0.0008                      | 0.0019   |
| 1997 | 0.0025                                               | 0.0023                     | 0.0010                      | 0.0021   |
| 1998 | 0.0024                                               | 0.0026                     | 0.0015                      | 0.0023   |
| 1999 | 0.0029                                               | 0.0023                     | 0.0016                      | 0.0024   |
| 2000 | 0.0020                                               | 0.0028                     | 0.0013                      | 0.0021   |
| 2001 | 0.0029                                               | 0.0023                     | 0.0018                      | 0.0024   |
| 2002 | 0.0019                                               | 0.0013                     | 0.0020                      | 0.0016   |
| 2003 | 0.0015                                               | 0.0016                     | 0.0017                      | 0.0016   |
| 2004 | 0.0021                                               | 0.0015                     | 0.0009                      | 0.0016   |
| 2005 | 0.0022                                               | 0.0013                     | 0.0006                      | 0.0015   |
| 2006 | 0.0019                                               | 0.0009                     | 0.0003                      | 0.0012   |
| 2007 | 0.0014                                               | 0.0009                     | 0.0004                      | 0.0010   |
| 2008 | 0.0014                                               | 0.0012                     | 0.0003                      | 0.0011   |
| 2009 | 0.0016                                               | 0.0014                     | 0.0004                      | 0.0012   |
| 2010 | 0.0011                                               | 0.0011                     | 0.0003                      | 0.0009   |
| 2011 | 0.0013                                               | 0.0009                     | 0.0004                      | 0.0009   |
| 2012 | 0.0010                                               | 0.0006                     | 0.0005                      | 0.0007   |
| 2013 | 0.0010                                               | 0.0009                     | 0.0005                      | 0.0008   |
| 2014 | 0.0007                                               | 0.0011                     | 0.0005                      | 0.0008   |

Annual upper 95% CIs for  $F$  in each region and the assessment area (three regions combined) (rounded to four decimal places).

| Year | Western<br>Indian<br>Ocean/Eastern<br>Atlantic Ocean | Eastern<br>Indian<br>Ocean | Western<br>Pacific<br>Ocean | Combined |
|------|------------------------------------------------------|----------------------------|-----------------------------|----------|
| 1992 | 0.0007                                               | 0.0007                     | 0.0005                      | 0.0006   |
| 1993 | 0.0010                                               | 0.0003                     | 0.0004                      | 0.0006   |
| 1994 | 0.0007                                               | 0.0003                     | 0.0003                      | 0.0005   |
| 1995 | 0.0007                                               | 0.0004                     | 0.0002                      | 0.0005   |
| 1996 | 0.0006                                               | 0.0005                     | 0.0002                      | 0.0005   |
| 1997 | 0.0006                                               | 0.0006                     | 0.0002                      | 0.0005   |
| 1998 | 0.0006                                               | 0.0006                     | 0.0004                      | 0.0006   |
| 1999 | 0.0007                                               | 0.0006                     | 0.0004                      | 0.0006   |
| 2000 | 0.0005                                               | 0.0007                     | 0.0003                      | 0.0005   |
| 2001 | 0.0007                                               | 0.0006                     | 0.0005                      | 0.0006   |
| 2002 | 0.0005                                               | 0.0003                     | 0.0005                      | 0.0004   |
| 2003 | 0.0004                                               | 0.0004                     | 0.0004                      | 0.0004   |
| 2004 | 0.0005                                               | 0.0004                     | 0.0002                      | 0.0004   |
| 2005 | 0.0005                                               | 0.0003                     | 0.0002                      | 0.0004   |
| 2006 | 0.0005                                               | 0.0002                     | 0.0001                      | 0.0003   |
| 2007 | 0.0003                                               | 0.0002                     | 0.0001                      | 0.0002   |
| 2008 | 0.0003                                               | 0.0003                     | 0.0001                      | 0.0003   |
| 2009 | 0.0004                                               | 0.0003                     | 0.0001                      | 0.0003   |
| 2010 | 0.0003                                               | 0.0003                     | 0.0001                      | 0.0002   |
| 2011 | 0.0003                                               | 0.0002                     | 0.0001                      | 0.0002   |
| 2012 | 0.0002                                               | 0.0001                     | 0.0001                      | 0.0002   |
| 2013 | 0.0002                                               | 0.0002                     | 0.0001                      | 0.0002   |
| 2014 | 0.0002                                               | 0.0003                     | 0.0001                      | 0.0002   |

Annual lower 95% CIs for  $F$  in each region and the assessment area (three regions combined) (rounded to four decimal places).

| Year | Western<br>Indian<br>Ocean/Eastern<br>Atlantic Ocean | Eastern<br>Indian<br>Ocean | Western<br>Pacific<br>Ocean | Combined |
|------|------------------------------------------------------|----------------------------|-----------------------------|----------|
| 1992 | 0.0203                                               | 0.0213                     | 0.0153                      | 0.0196   |
| 1993 | 0.0309                                               | 0.0084                     | 0.0124                      | 0.0181   |
| 1994 | 0.0217                                               | 0.0104                     | 0.0086                      | 0.0145   |
| 1995 | 0.0212                                               | 0.0126                     | 0.0073                      | 0.0149   |
| 1996 | 0.0173                                               | 0.0170                     | 0.0058                      | 0.0147   |
| 1997 | 0.0193                                               | 0.0179                     | 0.0077                      | 0.0162   |
| 1998 | 0.0186                                               | 0.0200                     | 0.0113                      | 0.0176   |
| 1999 | 0.0225                                               | 0.0177                     | 0.0123                      | 0.0184   |
| 2000 | 0.0154                                               | 0.0215                     | 0.0096                      | 0.0165   |
| 2001 | 0.0220                                               | 0.0177                     | 0.0142                      | 0.0185   |
| 2002 | 0.0143                                               | 0.0101                     | 0.0150                      | 0.0126   |
| 2003 | 0.0119                                               | 0.0126                     | 0.0129                      | 0.0122   |
| 2004 | 0.0164                                               | 0.0118                     | 0.0071                      | 0.0126   |
| 2005 | 0.0167                                               | 0.0099                     | 0.0048                      | 0.0115   |
| 2006 | 0.0142                                               | 0.0069                     | 0.0026                      | 0.0089   |
| 2007 | 0.0105                                               | 0.0072                     | 0.0028                      | 0.0076   |
| 2008 | 0.0105                                               | 0.0091                     | 0.0025                      | 0.0083   |
| 2009 | 0.0120                                               | 0.0104                     | 0.0030                      | 0.0095   |
| 2010 | 0.0087                                               | 0.0082                     | 0.0025                      | 0.0072   |
| 2011 | 0.0097                                               | 0.0066                     | 0.0031                      | 0.0071   |
| 2012 | 0.0077                                               | 0.0044                     | 0.0036                      | 0.0055   |
| 2013 | 0.0074                                               | 0.0066                     | 0.0039                      | 0.0063   |
| 2014 | 0.0051                                               | 0.0086                     | 0.0040                      | 0.0062   |

## Appendix C *F*-ratios

Annual upper 95% CIs for *F*-ratio, the impact of pelagic longline fisheries on porbeagle shark in each region and the assessment area (three regions combined) (rounded to three decimal places).

| Year | Western Indian Ocean/Eastern Atlantic Ocean | Eastern Indian Ocean | Western Pacific Ocean | Combined |
|------|---------------------------------------------|----------------------|-----------------------|----------|
| 1992 | 0.828                                       | 0.882                | 0.634                 | 0.906    |
| 1993 | 1.345                                       | 0.370                | 0.517                 | 0.731    |
| 1994 | 0.923                                       | 0.437                | 0.327                 | 0.596    |
| 1995 | 0.904                                       | 0.545                | 0.298                 | 0.602    |
| 1996 | 0.721                                       | 0.711                | 0.250                 | 0.596    |
| 1997 | 0.822                                       | 0.774                | 0.323                 | 0.721    |
| 1998 | 0.769                                       | 0.817                | 0.503                 | 0.738    |
| 1999 | 0.916                                       | 0.714                | 0.496                 | 0.734    |
| 2000 | 0.678                                       | 0.944                | 0.405                 | 0.710    |
| 2001 | 0.994                                       | 0.784                | 0.598                 | 0.784    |
| 2002 | 0.579                                       | 0.409                | 0.622                 | 0.474    |
| 2003 | 0.498                                       | 0.525                | 0.522                 | 0.487    |
| 2004 | 0.696                                       | 0.541                | 0.287                 | 0.526    |
| 2005 | 0.685                                       | 0.431                | 0.216                 | 0.509    |
| 2006 | 0.602                                       | 0.298                | 0.110                 | 0.361    |
| 2007 | 0.460                                       | 0.303                | 0.117                 | 0.307    |
| 2008 | 0.436                                       | 0.384                | 0.104                 | 0.335    |
| 2009 | 0.497                                       | 0.456                | 0.126                 | 0.392    |
| 2010 | 0.375                                       | 0.373                | 0.098                 | 0.325    |
| 2011 | 0.400                                       | 0.272                | 0.131                 | 0.278    |
| 2012 | 0.310                                       | 0.178                | 0.150                 | 0.244    |
| 2013 | 0.285                                       | 0.272                | 0.164                 | 0.264    |
| 2014 | 0.204                                       | 0.356                | 0.176                 | 0.245    |

Annual lower 95% CIs for *F*-ratio, the impact of pelagic longline fisheries on porbeagle shark in each region and the assessment area (three regions combined) (rounded to three decimal places).

| Year | Western Indian Ocean/Eastern Atlantic Ocean | Eastern Indian Ocean | Western Pacific Ocean | Combined |
|------|---------------------------------------------|----------------------|-----------------------|----------|
| 1992 | 0.014                                       | 0.015                | 0.011                 | 0.014    |
| 1993 | 0.021                                       | 0.006                | 0.009                 | 0.012    |
| 1994 | 0.014                                       | 0.007                | 0.006                 | 0.010    |
| 1995 | 0.014                                       | 0.008                | 0.005                 | 0.011    |
| 1996 | 0.012                                       | 0.012                | 0.004                 | 0.010    |
| 1997 | 0.014                                       | 0.013                | 0.005                 | 0.011    |
| 1998 | 0.013                                       | 0.014                | 0.008                 | 0.012    |
| 1999 | 0.016                                       | 0.012                | 0.008                 | 0.013    |
| 2000 | 0.011                                       | 0.014                | 0.007                 | 0.011    |
| 2001 | 0.015                                       | 0.012                | 0.010                 | 0.014    |
| 2002 | 0.010                                       | 0.007                | 0.010                 | 0.009    |
| 2003 | 0.008                                       | 0.009                | 0.009                 | 0.008    |
| 2004 | 0.011                                       | 0.008                | 0.005                 | 0.008    |
| 2005 | 0.011                                       | 0.007                | 0.003                 | 0.008    |
| 2006 | 0.010                                       | 0.005                | 0.002                 | 0.006    |
| 2007 | 0.007                                       | 0.005                | 0.002                 | 0.005    |
| 2008 | 0.007                                       | 0.006                | 0.002                 | 0.006    |
| 2009 | 0.008                                       | 0.007                | 0.002                 | 0.007    |
| 2010 | 0.006                                       | 0.006                | 0.002                 | 0.005    |
| 2011 | 0.007                                       | 0.005                | 0.002                 | 0.005    |
| 2012 | 0.005                                       | 0.003                | 0.002                 | 0.004    |
| 2013 | 0.005                                       | 0.004                | 0.003                 | 0.004    |
| 2014 | 0.003                                       | 0.006                | 0.003                 | 0.004    |

# Addendum

The original report chose one MIST (which is the minimum unsustainable F); all the results in the main body of the report are based on this one MIST. The WCPFC has considered several available options for the MIST<sup>1</sup> but has not yet provided guidance on which MIST to use. We have therefore decided to cater for differing definitions of applicable risk, and provide results against  $F_{msm}$  and  $F_{lim}$  MISTs as well as  $F_{crash}$ . Feedback on this addendum and the appropriate MIST can be incorporated into the main report when it is finalized after SC13.

Please note that the figures below are labelled ADD1, ADD2, and ADD3, corresponding to and supplementing Figures 21, 22 and 23 in the main body of the report.

---

<sup>1</sup> Clarke, S. and S. Hoyle. 2014. Development of limit reference points for elasmobranchs. WCPFC-SC10-2014/MI-WP-07. Accessed online at <https://www.wcpfc.int/system/files/SC10-MI-WP-07Development%20of%20LRPs%20for%20Elasmobranchs%20%28rev%20final%29.pdf>






Figure ADD1: Estimated fishing mortalities by year and region. The horizontal dotted lines indicate the median MIST values ( $F_{crash}$ ,  $F_{lim}$ , and  $F_{msm}$ ), but note that the MIST varies among realisations of the model, with the grey band showing the 90% distribution for  $F_{crash}$ . The boxes show the interquartile range of the  $F$  estimates, with a line at the median of each. The whiskers extend up to 1.5x the interquartile range. Dots mark points beyond the whiskers.




Figure ADD2: F-ratio plots showing the median values of  $F / MIST$  by year, for the three versions of the MIST ( $F_{msm}$ ,  $F_{im}$ , and  $F_{crash}$ ) for each of the three regions and for the three regions combined (the assessment area). Note that the F-ratio is almost always below 1, indicated by the horizontal dotted line.




Figure ADD3: Risk plots showing the probability that  $F$  exceeds the MIST by year, for the three versions of the MIST ( $F_{crash}$ ,  $F_{lim}$ , and  $F_{crash}$ ) for each of the three regions and for the three regions combined (the assessment area).