

EXECUTIVE SUMMARY: BLACK MARLIN

Status of the Indian Ocean black marlin (BLM: *Makaira indica*) resourceTABLE 1. Black marlin: Status of black marlin (*Makaira indica*) in the Indian Ocean.

Area ¹	Indicators		2017 stock status determination
Indian Ocean	Catch 2016 ² :	17,829 t	80%*
	Average catch 2012–2016:	16,638 t	
	MSY (1,000 t) (80% CI):	9.932 (6.963-12.153)	
	F _{MSY} (80% CI):	0.211 (0.089-0.430)	
	B _{MSY} (1,000 t) (80% CI):	47.430 (27.435-100.109)	
	F ₂₀₁₅ /F _{MSY} (80% CI):	2.42 (1.52-4.06)	
	B ₂₀₁₅ /B _{MSY} (80% CI):	0.81 (0.55-1.10)	
	B ₂₀₁₅ /B ₀ (80% CI):	0.30 (0.20-0.41)	

¹Boundaries for the Indian Ocean = IOTC area of competence;

²Proportion of catch estimated or partially estimated by IOTC Secretariat in 2016: 42%

* Estimated probability that the stock is in the respective quadrant of the Kobe plot (shown below), derived from the confidence intervals associated with the current stock status.

Colour key	Stock overfished (B _{year} /B _{MSY} < 1)	Stock not overfished (B _{year} /B _{MSY} ≥ 1)
Stock subject to overfishing (F _{year} /F _{MSY} > 1)	80%	19%
Stock not subject to overfishing (F _{year} /F _{MSY} ≤ 1)	0%	1%
Not assessed/Uncertain		

INDIAN OCEAN STOCK – MANAGEMENT ADVICE

Stock status. Stock status based on BSP-SS stock assessment suggests that the stock in 2015 is in the red zone in the Kobe plot with F/F_{MSY}=2.42 and B/B_{MSY}=0.81. Another approach by ASPIC examined in 2016 came to similar conclusions. The Kobe plot (Fig. 2) from the BSP-SS model indicated that the stock has been **subject to overfishing** and **overfished** in recent years (Table 1; Fig. 2).

Outlook. The uncertainty in the data available for assessment purposes and the CPUE series suggests that the advice should be interpreted with caution. The recent sharp increase of catch changed the status of stock to the red zone of the Kobe plot. Even if the catch levels are reduced by 40% of the average 2013-2015 levels, it is unlikely that biomass would be above the B_{MSY} and F would be below the F_{MSY} in the next 10 years (Table 2).

Management advice. The current catches (Fig. 1) are considerably higher than MSY (9,932 t) and the stock is overfished (B₂₀₁₅ < B_{MSY}) and currently subject to overfishing (F₂₀₁₅ > F_{MSY}). Even with a 40% reduction in current catches, it is very unlikely (less than 5%) to achieve the Commission objectives of being in the green zone of the Kobe Plot by 2025. Current catch levels are not sustainable and there is a need for urgent actions to decrease these catch levels. In order to enable the stock to start rebuilding, the Commission should consider a reduction of substantially greater than 40% from the current catches.

The following key points should be noted:

- **Maximum Sustainable Yield (MSY):** estimate for the whole Indian Ocean is 9,932 t.
- **Provisional reference points:** Although the Commission adopted reference points for swordfish in Resolution 15/10 *on target and limit reference points and a decision framework*, no such interim reference points nor harvest control rules have been established for black marlin.
- **Main fishing gear (average catches 2012–16):** Black marlin are largely considered to be a non-target species of industrial and artisanal fisheries. Gillnets account for around 53% of total catches in the Indian Ocean, followed by longlines (17%), with remaining catches recorded under troll and handlines. (Fig. 1).
- **Main fleets (average catches 2012-16):** Iran (≈28%), India (≈20%), Sri Lanka (≈20%), and Indonesia (≈15%).





Fig. 1. Black marlin: catches by gear and year recorded in the IOTC Database (1950–2016)¹.



¹ **Definition of fisheries:** Longline; Gillnet; Hook-and-Line (includes handline, trolling, baitboat, and sport fisheries); Other gears (includes coastal purse seine, Danish purse seine, beach seine, and purse seine).

Fig. 2. Black marlin: BSP-SS aggregated Indian Ocean assessment Kobe plots for black marlin (contours are the 25, 50, 75 and 90 percentiles of the 2015 estimate). Black line indicates the trajectory of the point estimates for the total biomass (B) ratio and F ratio for each year 1950–2015.

Table 2. Black Marlin: Indian Ocean BSP-SS Kobe II Strategy Matrix. Probability (percentage) of violating the MSY-based reference points for nine constant catch projections (average catch level from 2013 to 2015 17,171 t, $\pm 10\%$, $\pm 20\%$, $\pm 30\%$ $\pm 40\%$) projected for 3 and 10 years.

Reference point and projection timeframe	Alternative catch projections (relative to the average catch level from 2013 to 2015, 17,171 t, and probability (%) of violating MSY-based reference points								
	60%	70%	80%	90%	100%	110%	120%	130%	140%
	10,303 t	12,020 t	13,737 t	15,454 t	17,171 t	18,888 t	20,605 t	22,322 t	24,039 t
B ₂₀₁₈ < B _{MSY}	91	94	96	97	98	98	99	99	99
F ₂₀₁₈ > F _{MSY}	89	96	98	99	100	100	100	100	100
B ₂₀₂₅ < B _{MSY}	98	100	100	100	100	100	100	100	100
F ₂₀₂₅ > F _{MSY}	97	99	100	100	100	100	100	100	100