

OUTCOMES OF THE FOURTEENTH SESSION OF THE SCIENTIFIC COMMITTEE

PREPARED BY: IOTC SECRETARIAT, 15 JUNE, 2012

PURPOSE

To inform the Working Party on Temperate Tunas (WPTmT) of the recommendations arising from the Fourteenth Session of the Scientific Committee, held from 12–17 December 2011, specifically relating to the work of the WPTmT.

BACKGROUND

At the Fourteenth Session of the Scientific Committee (SC), the recommendations relevant to the work of the WPTmT contained in [Appendix A](#) were adopted by the SC and provided to the Commission for its consideration.

In addition, the SC noted and endorsed the recommendations made by the WPTmT in 2011, which included requests to address the deficiencies in data collection, monitoring and reporting by CPCs. The SC requested that the IOTC Secretariat communicate these recommendations to relevant parties so that they may address these matters in 2012 and provide progress updates to the WPTmT at its next meeting.

The recommendations on the deficiencies in data collection, monitoring and reporting by CPCs in relation to temperate tunas will be discussed under agenda item 5 and in paper IOTC-2012-WPTmT04-06 and are therefore not presented in this paper.

DISCUSSION

In addition to the recommendations outlined in [Appendix A](#), the SC made several other comments and recommendations relevant to the WPTmT, which participants are asked to consider:

- The SC **NOTED** that the catches of albacore estimated for the fresh tuna longline fishery of Indonesia in recent years are thought to be uncertain, as they cannot be verified using data collected through port sampling, and that to date, the IOTC Secretariat has not received catch-and-effort data for this fishery. The SC was also informed that misidentification between yellowfin tuna and albacore might occur in the Indonesian catches which may contribute to the rise of declared albacore catches in recent years. However, the catch levels estimated by the IOTC Secretariat also account for other sources such as the export declarations from Bali and canning factories receiving the products abroad. Finally, the SC urged Indonesia to undertake a thorough examination of the sampling procedure at landing sites as soon as possible. Indonesia requested that the IOTC Secretariat to bridge the gap of catch data of albacore recorded by Indonesian authorities by providing a list of vessels directly exporting albacore to the canning factories abroad. (para. 28 of the SC14 report)
- The SC **NOTED** the difficulties faced by Indonesian scientists and managers in terms of commercial catches being transhipped at sea, as well as catches directly exported abroad contributing to IUU fishing. The SC **HIGHLIGHTED** the need for logbooks to be utilised on all commercial fishing vessels, noting that this is already a mandatory requirement for IOTC CPCs. Indonesia encouraged collaboration among CPCs to exchange necessary information related to vessels landing their catch to their countries. (para. 29 of the SC14 report)

Implementation of the regional observer scheme

- The SC **NOTED** the update on the implementation of the Regional Observer Scheme set out in Resolution 11/06 *on a Regional Observer Scheme* and **EXPRESSED** its concerns regarding the low level of implementation and reporting to the IOTC Secretariat of both the observer trip reports and the list of

accredited observers since the start of the ROS in July 2010 (8 CPCs provided a list of accredited observers and 11 reports were submitted from 4 CPCs). (para. 138 of the SC14 report)

- The SC **RECOMMENDED** that all IOTC CPCs urgently implement the requirements of Resolution 11/04 on a Regional Observer Scheme, which states that: The observer shall, within 30 days of completion of each trip, provide a report to the CPCs of the vessel. The CPCs shall send within 150 days at the latest each report, as far as continuous flow of report from observer placed on the longline fleet is ensured, which is recommended to be provided with 1°x1° format to the Executive Secretary, who shall make the report available to the Scientific Committee upon request. In a case where the vessel is fishing in the EEZ of a coastal state, the report shall equally be submitted to that Coastal State. (para. 11), **NOTING** that the timely submission of observer trip reports to the Secretariat is necessary to ensure that the Scientific Committee is able to carry out the tasks assigned to it by the Commission, including the analysis of accurate and high resolution data, in particular for bycatch, which would allow the scientists to better assess the impacts of fisheries for tuna and tuna-like species on bycatch species. (para. 139 of the SC14 report)

Management strategy evaluation

- The SC **ENDORSED** the roadmap presented for the implementation of MSE in the Indian Ocean in IOTC-2011-SC14-36 and **RECOMMENDED** the Commission agree to initiate a consultative process among managers, stakeholders and scientists to begin discussions about the implementation of MSE in IOTC. (para. 157 of the SC14 report)

RECOMMENDATION

That the WPTmT **NOTE** the recommendations of the Fourteenth Session of the Scientific Committee and consider how to progress these issues at the present meeting.

APPENDICES

Appendix A: Consolidated set of recommendations of the Fourteenth Session of the Scientific Committee (12–17 December, 2011) to the Commission, relevant to the Working Party on Temperate Tunas.

Appendix B: Status of the Indian Ocean albacore tuna resource.

APPENDIX A

**CONSOLIDATED SET OF RECOMMENDATIONS OF THE FOURTEENTH SESSION OF THE
SCIENTIFIC COMMITTEE (12–17 DECEMBER, 2011) TO THE COMMISSION RELEVANT TO
WORKING PARTY ON TEMPERATE TUNAS**

Extract of the Report of the Fourteenth Session of the Scientific Committee

(IOTC-2011-SC14-R; Appendix XXXVIII, PAGES 248–259)

STATUS OF TUNA AND TUNA-LIKE RESOURCES IN THE INDIAN OCEAN

Tuna – Highly migratory species

- SC14.01 (para. 129) The SC **RECOMMENDED** that the Commission note the management advice developed for each tropical and temperate tuna species as provided in the Executive Summary for each species.
- Albacore (*Thunnus alalunga*) – [Appendix X](#) [provided as Attachment B below]

GENERAL RECOMMENDATIONS TO THE COMMISSION

Activities of the IOTC Secretariat in 2011

- SC14.10 (para. 32) Noting the request by the Commission at its 15th Session for a new assessment of albacore to be undertaken in 2011 (para. 37 of the S15 report), the SC **RECOMMENDED** that the Commission note that although a new assessment was undertaken in 2011, there remains considerable uncertainty about the relationship between abundance and the standardized CPUE series, and about the total catches over the past decade and that the WPTmT has limited confidence in the assessment undertaken. Thus, there is an urgent need to carry out a revised stock assessment for the albacore resource in the Indian Ocean in 2012, and the Commission should consider allocating funds for this purpose, noting that individual CPCs are finding it difficult to justify expending the necessary resources to undertake stock assessments.

Examination of the Effect of Piracy on Fleet Operations and Subsequent Catch and Effort Trends

- SC14.46 (para. 127) In response to the request of the Commission (para. 40 of the S15 report), the SC **RECOMMENDED** that given the lack of quantitative analysis of the effects of piracy on fleet operations and subsequent catch and effort trends, and the potential impacts of piracy on fisheries in other areas of the Indian Ocean through the relocation of longliners to other fishing grounds, specific analysis should be carried out and presented at the next WPTT meeting by the CPCs most affected by these activities, including Japan, Republic of Korea and Taiwan, China.

Implementation of the Precautionary approach and Management strategy Evaluation

- SC14.49 (para. 146) Noting that the development of an MSE process will require management objectives to be specified, the SC **RECOMMENDED** that the Commission provide clear guidance in this regard, noting that the adoption of the Precautionary Approach, as defined in the Fish Stocks Agreement, may be the first step.

- SC14.50 (para. 149) The SC **RECOMMENDED** that interim target and limit reference points be adopted and a list of possible provisional values for the major species is listed in [Table 5](#). These values should be replaced as soon as the MSE process is completed. Provisional target reference points would be based on the MSY level of the indicators, and on different multipliers for the limit reference points.

Table 5. Interim target and limit reference points.

Stock	Target Reference Point	Limit Reference Point
Albacore	$B_{MSY}; F_{MSY}$	$0.4*B_{MSY}; 1.4*F_{MSY}$
Bigeye tuna	$B_{MSY}; F_{MSY}$	$0.5*B_{MSY}; 1.3*F_{MSY}$
Skipjack tuna	$B_{MSY}; F_{MSY}$	$0.4*B_{MSY}; 1.5*F_{MSY}$
Yellowfin tuna	$B_{MSY}; F_{MSY}$	$0.4*B_{MSY}; 1.4*F_{MSY}$
Swordfish	$B_{MSY}; F_{MSY}$	$0.4*B_{MSY}; 1.4*F_{MSY}$

Evaluation of the IOTC time-area closure

- SC14.57 (para. 178) The SC **RECOMMENDED** that the Commission note that the current closure is likely to be ineffective, as fishing effort will be redirected to other fishing grounds in the Indian Ocean. The positive impacts of the moratorium within the closed area would likely be offset by effort reallocation. For example, the WPTmT noted that longline fishing effort has been redistributed to traditional albacore fishing grounds in recent years, thereby further increasing fishing pressure on this stock.
- SC14.58 (para. 179) Noting that the objective of Resolution 10/01 is to decrease the overall pressure on the main targeted stocks in the Indian Ocean, in particular yellowfin tuna and bigeye tuna, and also to evaluate the impact of the current time/area closure and any alternative scenarios on tropical tuna population, the SC **RECOMMENDED** that the Commission specify the level of reduction or the long term management objectives to be achieved with the current or alternative time area closures, as these are not contained within the Resolution 10/01.

RESEARCH RECOMMENDATIONS AND PRIORITIES

Stock assessment


- SC14.71 (para. 206) The SC **AGREED** that there was an urgent need to carry out revised stock assessments for the albacore resource in the Indian Ocean in 2012, and **RECOMMENDED** that the Commission consider approving funds for this purpose.

Stock structure

- SC14.72 (para. 207) Noting that at present very little is known about the population structure and migratory range of albacore in the Indian Ocean, other than the possible connectivity with the southern Atlantic, the SC **RECOMMENDED** that a research project addressing the albacore stock structure, migratory range and movement rates in the Indian Ocean be considered at its 2012 annual meeting as this project is assigned a high priority.

Additional core topics for research

- SC14.73 (para. 208) The SC **RECOMMENDED** that the following core topic areas as priorities for research over the coming year:
- Size data analyses
 - Growth rates and ageing studies
 - Stock status indicators – exploration of indicators from available data
 - Collaborate with SPC-OFP to examine their current simulation approach to determine priority research areas.



Status of the Indian Ocean Albacore Tuna Resource (*Thunnus alalunga*)

TABLE 1. Status of albacore (*Thunnus alalunga*) in the Indian Ocean.

Area ¹	Indicators – 2011 assessment		2011 stock status determination
			2010 ²
Indian Ocean	Catch 2010: 43,711 t Average catch 2006–2010: 41,074 t MSY (1 model): 29,900 t (21,500–33,100 t) F_{2010}/F_{MSY} (1 model): 1.61 (1.19–2.22)* B_{2010}/B_{MSY} (1 model): 0.89 (0.65–1.12)* B_{2010}/B_{1980} (1 model): 0.39 (n.a.)		

¹Boundaries for the Indian Ocean stock assessment are defined as the IOTC area of competence.

²The stock status refers to the most recent years' data used for the assessment.

*(Note: at this time the WPTmT had limited confidence in the assessment results (refer to paragraphs 71–77 in the report of the WPTmT03 (IOTC–2011–WPTmT03–R) for further clarification).

Colour key	Stock overfished ($SB_{year}/SB_{MSY} < 1$)	Stock not overfished ($SB_{year}/SB_{MSY} \geq 1$)
Stock subject to overfishing ($F_{year}/F_{MSY} > 1$)		
Stock not subject to overfishing ($F_{year}/F_{MSY} \leq 1$)		

INDIAN OCEAN STOCK – MANAGEMENT ADVICE

The SC **RECOMMENDED** the following management advice for albacore in the Indian Ocean noting that there remains considerable uncertainty about the relationship between abundance and the standardized CPUE series, and about the total catches over the past decade.

Stock status. Trends in the Taiwan, China CPUE series suggest that the longline vulnerable biomass has declined to about 39% of the level observed in 1980. There were 20 years of moderate fishing before 1980, and the catch has more than doubled since 1980. Catches have increased substantially since the previous albacore assessment when there was considered to be a risk that $SB < SB_{MSY}$, so the risk will have increased further. It is considered likely that recent catches have been above MSY, recent fishing mortality exceeds F_{MSY} ($F_{2010}/F_{MSY} > 1$). There is a moderate risk that total biomass is below B_{MSY} ($B_{2010}/B_{MSY} \approx 1$) (Table 1, Fig. 1).

Outlook. Maintaining or increasing effort will probably result in further declines in biomass, productivity and CPUE. The impacts of piracy in the western Indian Ocean has resulted in the displacement of a substantial portion of longline fishing effort into the traditional albacore fishing areas in the southern and eastern Indian Ocean. It is therefore unlikely that catch and effort on albacore will decline in the near future.

The SC **RECOMMENDED** the following:

- The available evidence indicates considerable risk to the stock status at current effort levels.
- The two primary sources of data that drive the assessment, total catches and CPUE are highly uncertain and should be investigated further as a priority.
- Current catches (average ~41,000 t over the last five years, ~44,000 t in 2010) likely exceed MSY (29,900 t, range: 21,500–33,100 t). Maintaining or increasing effort will probably result in further declines in biomass, productivity and CPUE.
- A Kobe 2 Strategy matrix was calculated to quantify the risk of different future catch scenarios. However, a number of inconsistencies between the model and data were noted for future investigation (matrix not presented here as a result).




Fig. 1. ASPIC Aggregated Indian Ocean assessment Kobe plot (95% Confidence surfaces shown around 2010 estimate). Fixed $B/K=0.9$. Blue circles indicate the trajectory of the point estimates for the TB ratio and F ratio for each year 1980–2010 (Note: at this time the WPTmT had limited confidence in the assessment results (refer to paragraphs 71–77 in the report of the WPTmT03 (IOTC–2011–WPTmT03–R) for further clarification).

SUPPORTING INFORMATION

(Information collated from reports of the Working Party on Temperate Tunas and other sources as cited)

CONSERVATION AND MANAGEMENT MEASURES

Albacore (*Thunnus alalunga*) in the Indian Ocean are currently subject to a number of conservation and management measures adopted by the Commission, although none are species specific:

- Resolution 08/04 concerning the recording of catch by longline fishing vessels in the IOTC area.
- Resolution 09/02 On the implementation of a limitation of fishing capacity of contracting parties and cooperating non-contracting parties.
- Resolution 10/02 mandatory statistical requirements for IOTC Members and Cooperating non-Contracting Parties (CPC's).
- Resolution 10/03 concerning the recording of catch by fishing vessels in the IOTC area.
- Resolution 10/07 concerning a record of licensed foreign vessels fishing for tunas and swordfish in the IOTC area.
- Resolution 10/08 concerning a record of active vessels fishing for tunas and swordfish in the IOTC area.
- Recommendation 11/06 Concerning the Recording of Catch by Fishing Vessels in the IOTC Area of Competence.

FISHERIES INDICATORS

General

Overall, the biology of albacore stock in the Indian Ocean is not well known and there is relatively little new information on albacore stocks. Albacore (*Thunnus alalunga*) life history characteristics, including a relatively late maturity, long life and sexual dimorphism, make the species vulnerable to over exploitation. Table 2 outlines some of the key life history traits of albacore specific to the Indian Ocean.

Catch trends

Albacore are currently caught almost exclusively using drifting longlines (98%), and between 20°S and 40°S, with remaining catches recorded using purse seines and other gears (Fig. 2). Between 1983 and 1992, a large portion of albacore catches were taken by the Taiwan, China fleet using drifting gillnets (Fig. 2; Table 3) which targeted juvenile albacore in the southern Indian Ocean (30°S to 40°S). In 1992 the United Nations worldwide ban on the use of drifting gillnets effectively closed this gillnet fishery.

Catches of albacore were relatively stable until the mid-1980s, except for high catches recorded in 1973 and 1974 (Fig. 2). The catches increased markedly during the mid-1980's due to the use of drifting gillnets by Taiwan, China, with total catches in excess of 30,000 t. Following the removal of the drifting gillnet fleet, catches dropped to less than 20,000 t by 1993. However, catches more than doubled over the period from 1993 (less than 20,000 t) to 2001 (44,000 t). Record catches of albacore were reported in 2007, at around 45,000 t, and again in 2008, at 48,000 t. Catches for 2009 are estimated to be approximately 40,000 t, while preliminary catches for 2010 amount to 43,711 t (Table 3).

TABLE 2. Biology of Indian Ocean albacore (*Thunnus alalunga*)

Parameter	Description
Range and stock structure	<p>A temperate tuna living mainly in the mid oceanic gyres of the Pacific, Indian and Atlantic oceans. In the Pacific and Atlantic oceans there is a clear separation of southern and northern stocks associated with the oceanic gyres that are typical of these areas. In the Indian Ocean, there is probably only one southern stock, distributed from 5°N to 40°S, because there is no northern gyre.</p> <p>Albacore is a highly migratory species and individuals swim large distances during their lifetime. It can do this because it is capable of thermoregulation, has a high metabolic rate, and advanced cardiovascular and blood/gas exchange systems. Pre-adults (2-5 year old albacore) appear to be more migratory than adults. In the Pacific Ocean, the migration, distribution availability, and vulnerability of albacore are strongly influenced by oceanographic conditions, especially oceanic fronts. It has been observed on all albacore stocks that juveniles concentrate in cold temperate areas (for instance in a range of sea-surface temperatures between 15 and 18°C), and this has been confirmed in the Indian Ocean where albacore tuna are more abundant north of the subtropical convergence (an area where these juvenile were heavily fished by driftnet fisheries during the late 1980's). It appears that juvenile albacore show a continuous geographical distribution in the Atlantic and Indian oceans in the north edge of the subtropical convergence. Albacore may move across the jurisdictional boundary between ICCAT and IOTC.</p> <p>It is likely that the adult Indian Ocean albacore tunas do yearly circular counter-clockwise migrations following the surface currents of the south tropical gyre between their tropical spawning and southern feeding zones. In the Atlantic Ocean, large numbers of juvenile albacore are caught by the South African pole-and-line fishery (catching about 10,000 t yearly) and it has been hypothesized that these juveniles may be taken from a mixture of fish born in the Atlantic (north east of Brazil) and from the Indian Ocean. For the purposes of stock assessments, one pan-ocean stock has been assumed.</p>
Longevity	8 years (reported to 10 years in the Pacific)
Maturity (50%)	<p>Age: females 5–6 years; males n.a.</p> <p>Size: females n.a.; males n.a.</p>
Spawning season	Little is known about the reproductive biology of albacore in the Indian Ocean but it appears, based on biological studies and on fishery data, that the main spawning grounds are located east of Madagascar between 15° and 25°S during the 4th and 1st quarters of each year. Like other tunas, adult albacore spawn in warm waters (SST>25°C).
Size (length and weight)	n.a.

n.a. = not available. SOURCES: Froese & Pauly (2009); Xu & Tian (2011)

Catches of albacore in recent years have come almost exclusively from vessels flagged in Indonesia and Taiwan,China, although the catches of albacore reported for the fresh tuna longline fishery of Indonesia have increased considerably since 2003 to around 17,000 t (Fig. 3), which represents approximately 40% of the total catches of albacore in the Indian Ocean.

Longliners from Japan and Taiwan,China have been operating in the Indian Ocean since the early 1950s (Fig. 3). While the Japanese albacore catch ranged from 8,000 t to 18,000 t in the period 1959 to 1969, in 1972 catches rapidly decreased to around 1,000 t, due to a change in the target species, mainly to southern bluefin tuna and bigeye tuna. Albacore became a bycatch species for the Japanese fleet with catches between 200 t and 2,500 t. In recent years the Japanese albacore catch has been around 2,000 to 6,000 t.

In contrast to the Japanese longliners, catches by Taiwan,China longliners increased steadily from the 1950's to average around 10,000 t by the mid-1970s. Between 1998 and 2002 catches ranged between 21,500 t to 26,900 t, equating to just over 60% of the total Indian Ocean albacore catch. Between 2003 and 2010 the albacore catches by Taiwan,China longliners have been between 10,000 and 18,000 t, with catches appearing to be on the increase in recent years. There has been a shift in the proportion of catches of albacore by deep-freezing and fresh-tuna longliners in recent years, with increasing catches of fresh-tuna (68% of the total catches for 2008–2010) as opposed to deep-freezing longliners (Fig. 2; Table 3).

While most of the catches of albacore have traditionally come from the western Indian Ocean, in recent years a larger proportion of the catch has come from the southern and eastern Indian Ocean (Fig. 4; Table 4). The relative increase in catches in the eastern Indian Ocean since the early 2000's is mostly due to increased activity of fresh-tuna longliners from Taiwan,China and Indonesia (Indonesia not represented in Fig. 4 as spatial catch-and-effort data is not available or highly uncertain for these fleets). In the western Indian Ocean, the catches of albacore mostly result from the activities of deep-freezing longliners and purse seiners.

Fleets of oceanic gillnet vessels from Iran and Pakistan and gillnet and longline vessels from Sri Lanka have extended their area of operation in recent years, to operate on the high seas closer to the equator. The lack of catch-and-effort data from these fleets makes it impossible to assess whether they are operating in areas where catches of juvenile albacore are likely to occur.




Fig. 2. Annual catches of albacore by gear recorded in the IOTC Database (1961–2010) (Data as of August 2011). Other gears nei (Other); Purse seine (PS); Freezing-longline (LL); Fresh-tuna longline (FLL); Swordfish-longline (ELL).




Fig. 3. Annual catches of albacore by fleet recorded in the IOTC Database (1961–2010) (Data as of August 2011).





Fig. 4a–b. Time-area catches (total combined in tonnes) of albacore estimated for 2009 (left) and 2010 (right) by type of gear: Longline (LL, green), Driftnet (DFRT, red), Purse seine (PS, purple), Other fleets (OTHER, blue). Time-area catches are not available for all fleets; catches for those were assigned by 5x5 square and month using information from other fleets. Catches of fresh-tuna longliners are not represented (Data as of August 2011).

TABLE 3. Best scientific estimates of the catches of albacore (*Thunnus alalunga*) by gear and main fleets [or type of fishery] by decade (1950–2000) and year (2001–2010), in tonnes. Data as of October 2011. Catches by decade represent the average annual catch, noting that some gears were not used for all years (refer to Fig. 2).

Fishery	By decade (average)						By year (last ten years)									
	1950s	1960s	1970s	1980s	1990s	2000s	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
DN				5,823	3,735											
LL	3,715	17,231	16,900	15,212	21,876	20,283	38,664	29,998	17,818	16,283	16,149	14,123	11,468	11,704	12,874	14,498
FLL			80	314	1,329	15,493	3,728	3,920	6,910	15,242	15,524	14,455	31,759	33,969	26,619	28,752
FS				195	1,578	855	1,030	755	1,493	230	149	1,388	705	1,391	366	166
LS				8	105	65	251	17	3	2	15	160	21	33	26	42
OT	5	9	24	67	61	148	172	139	131	150	143	108	107	91	293	254
Total	3,721	17,240	17,005	21,620	28,684	36,844	43,845	34,829	26,355	31,906	31,979	30,234	44,059	47,189	40,178	43,711

Fisheries: Driftnet (DN); Taiwan, China); Freezing-longline (LL); Fresh-tuna longline (FLL); Purse seine free-school (FS); Purse seine associated school (LS); Other gears nei (OT). Note: LL includes the ELL catches shown in Fig. 2.

TABLE 4. Best scientific estimates of the catches of albacore (*Thunnus alalunga*) by fishing area for the period 1950–2009 (in metric tons). Data as of October 2011.

Area	By decade (average)						By year (last ten years)									
	1950s	1960s	1970s	1980s	1990s	2000s	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
N	3,369	8,275	7,659	5,486	8,525	21,597	20,526	13,867	14,049	19,538	19,809	18,625	34,248	30,189	29,827	23,257
S	351	8,965	9,346	16,134	20,158	15,247	23,319	20,962	12,306	12,368	12,170	11,609	9,811	17,000	10,351	20,454
Total	3,721	17,240	17,005	21,620	28,684	36,844	43,845	34,829	26,355	31,906	31,979	30,234	44,059	47,189	40,178	43,711

Areas: North of 10°S (N); South of 10°S (S)

Uncertainty of catches

Retained catches are fairly well known (Fig. 5); however catches are uncertain for:

- Longliners of Indonesia, India and Malaysia operating in Southern waters: To date, Indonesian, Indian and Malaysian longline vessels operating in Southern waters have not reported catches of albacore, noting that the Secretariat has estimated these catches at around 3000 t annually.
- Fleets using gillnets on the high seas, in particular Iran, Pakistan and Sri Lanka: Catches are likely to be less than 1000 t.
- Non-reporting industrial longliners (NEI): Refers to catches from longliners operating under flags of non-reporting countries. Historically high catches, however thought to be between 1000 and 2000 t in recent years.




Fig. 5. Uncertainty of annual catch estimates for albacore (1950–2010) (Data as of August 2011).

Catches below the zero-line (**Type B**) refer to fleets that do not report catch data to the IOTC (estimated by the IOTC Secretariat), do not report catch data by gear and/or species (broken by gear and species by the IOTC Secretariat) or any of the other reasons provided in the document. Catches over the zero-line (**Type A**) refer to fleets for which no major inconsistencies have been found to exist. Light bars represent data for artisanal fleets and dark bars represent data for industrial fleets.

- The catch series for albacore in recent years has changed substantially, especially since 2003. This change was due to a review of the data series for Indonesian longliners (Fig. 6).
- Levels of discards are believed to be low although they are unknown for industrial fisheries other than European (EU) purse seiners.
- Catch-and-effort series are available from various industrial fisheries. Nevertheless, catch-and-effort are not available from some fisheries or they are considered to be of poor quality, especially during the last decade, for the following reasons:
 - uncertain data from significant fleets of longliners, including India, Indonesia and Philippines.
 - non-reporting by industrial purse seiners and longliners (NEI).






Fig. 6. Catches of fresh-tuna longline vessels based in Indonesia (domestic and foreign) estimated in 2011 (1973–2010) versus catches estimated in 2008 (1973–2006). The revised Indonesian nominal catch series data was estimated by the IOTC Secretariat.

Effort trends

Total effort from longline vessels flagged to Japan, Taiwan, China and EU, Spain by five degree square grid from 2007 to 2010 are provided in Fig. 7, and total effort from purse seine vessels flagged to the EU and Seychelles (operating under flags of EU countries, Seychelles and other flags), and others, by five degree square grid and main fleets, for the years 2007 to 2010 are provided in Fig. 8.



Standardised catch-per-unit-effort (CPUE) trends

The CPUE series available for assessment purposes are shown in Fig. 9, although only the Taiwan,China series was used in the stock assessment model for 2011 for the reasons discussed in IOTC-2011-WPTmT03-R.




Fig. 9. Comparison of the three CPUE series for Indian Ocean albacore. Series have been rescaled relative to their respective means from 1960–2010.

Fish size or age trends (e.g. by length, weight, sex and/or maturity)

The size frequency data for the Taiwanese deep-freezing longline fishery for the period 1980–2009 is available. In general, the amount of catch for which size data for the species are available before 1980 is still very low. The data for the Japanese longline fleets is available; however, the number of specimens measured per stratum has been decreasing in recent years. Few data are available for the other fleets.

- Trends in average weight can be assessed for several industrial fisheries although they are incomplete or of poor quality for most fisheries before 1980, between 1986 and 1991, and in recent years, for the fleets referred to above (Fig. 10).
- Catch-at-Size(Age) tables are available but the estimates are highly uncertain for some periods and fisheries including:
 - all industrial longline fleets before the mid-60s, from the early-1970s up to the early-1980s and most fleets in recent years, in particular fresh-tuna longliners.
 - the paucity of catch by area data available for some industrial fleets (Taiwan,China, NEI, India and Indonesia).




Fig. 10. Average weight in kg of the catches of all fleets (blue), gillnet (red), LL-JPN (dark green), LL-TWN (black), Purseine (green) and other gears (grey) from 1950 to 2010.

STOCK ASSESSMENT

A single quantitative modelling method, a highly aggregated “A Stock Production Model Including Covariate” (ASPIC) surplus production model, was applied to the albacore assessment in 2011.

The following is worth noting with respect to the modelling approach used:

- The Taiwan,China CPUE standardisation should be used over the Japanese CPUE series because the Japanese CPUE demonstrates strong targeting shifts away from albacore (1960s) and toward albacore in recent years (as

a consequence of piracy in the western Indian Ocean), that was not accounted for in the standardization analysis.

- The Fox model had problems converging to a sensible solution when catch data prior to 1980 were included, when the Japanese CPUE were given substantial weight, and/or when the initial biomass was constrained to be less than or equal to the carrying capacity. The Working paper IOTC–2011–WPTmT03–19: *A note on the ASPIC Fox model and Indian Ocean albacore assessment*, examined this issue and found that the long catch time series tends to result in MSY estimates that approach 0. This causes a numerical failure. However, it appears that a range of MSY values may be reasonably consistent with the data.

The Fox model should be given a realistic biological constraint of $B(1980) < \text{carrying capacity}$ ($B(1980)/K=0.9$), otherwise the model estimates $B(1980) \gg K$. There was some incompatibility among the CPUE series, catch data and the Fox model. The structural rigidity of the Fox model limits the number of ways in which the error processes can be examined, and it was felt that this limited the scope of the analysis. Attempts to resolve the limitations are encouraged, as is the use of alternative models.

The general population trends and MSY parameters estimated by the Fox model appeared to be plausibly consistent with the general perception of the fishery and the data. However, these results are considered to be highly uncertain because of i) uncertainty in the catch rate standardization, ii) uncertainty in recent catches, and iii) limited ability to explore alternative interpretations of the data due to software constraints. The WPTmT had limited confidence in the assessment results.

TABLE 5. Albacore (*Thunnus alalunga*) stock status summary.

Management Quantity	Aggregate Indian Ocean
2010 catch estimate	43,700 t
Mean catch from 2006–2010	41,100 t
MSY (80% CI)	29,900 t (21,500–33,100)
Data period used in assessment	1980–2010
F_{2010}/F_{MSY} (80% CI)	1.61 (1.19–2.22)
B_{2010}/B_{MSY} (80% CI)	0.89 (0.65–1.12)
SB_{2010}/SB_{MSY}	–
B_{2010}/B_{1980} (80% CI)	0.39 (n.a.)
SB_{2010}/SB_{1980}	–
$B_{2010}/B_{1980, F=0}$	–
$SB_{2010}/SB_{1980, F=0}$	–

LITERATURE CITED

- Froese R, & Pauly DE 2009. *FishBase*, version 02/2009, FishBase Consortium, <www.fishbase.org>.
- Xu L & Tian SQ 2011. A study of fisheries biology for albacore based on Chinese observer data, IOTC–2011–WPTmT03–11.